

Testing tensor product Bézier surfaces for coincidence: A comprehensive solution

Krassimira Vlachkova^a

^aFaculty of Mathematics and Informatics, Sofia University "St. Kliment Ohridski", Blvd. James Bourchier 5, Sofia, 1164, Bulgaria

Abstract

It is known that Bézier curves and surfaces may have multiple representations by different control polygons. The polygons may have different number of control points and may even be disjoint. Up to our knowledge, Pekerman et al. (2005) were the first to address the problem of testing two parametric polynomial curves for coincidence. Their approach is based on reduction of the input curves into canonical irreducible form. They claimed that their approach can be extended for testing tensor product surfaces but gave no further detail.

In this paper we develop a new technique and provide a comprehensive solution to the problem of testing tensor product Bézier surfaces for coincidence. In (Vlachkova, 2017) an algorithm for testing Bézier curves was proposed based on subdivision. There a partial solution to the problem of testing tensor product Bézier surfaces was presented. Namely, the case where the irreducible surfaces are of same degree (n, m) , $n, m \in \mathbb{N}$, was resolved under certain additional condition. The other cases where one of the surfaces is of degree (n, m) and the other is of degree either $(n, n + m)$, or $(n + m, m)$, or $(n + m, n + m)$ remained open.

We have implemented our algorithm for testing tensor product Bézier surfaces for coincidence using Mathematica package. Experimental results and their analysis are presented.

Keywords: Bézier curve, Bézier surface, coincident surfaces, blossoming

1. Introduction

Assume we are given the control polygons of two tensor product Bézier (TPB) surfaces that are generated by different sources, e. g. different algorithms or software packages. The two polygons may have different number of control points and may even be disjoint but nevertheless it is possible that they represent surfaces with coincidence. Here we consider the problem of finding whether two control polygons represent different surfaces or partially/entirely coincident surfaces. In the latter case we need to determine their coincident part. By *coincident* surfaces we mean that they occupy same locus of points in \mathbb{R}^3 but they may be parameterized differently, i. e. they are *geometrically equivalent* as defined in (Denker and Herron, 1997).

This problem arises in various applications where the two surfaces need to be stitched together so that the obtained new surface is continuous. The problem is important also for the intersection algorithms based on subdivision which do not work well if the surfaces have coincident part.

The problem of testing polynomial curves for coincidence received a considerable attention by many authors. Up to our knowledge, Pekerman et al. (2005) were the first to address the problem of testing two parametric polynomial

*Corresponding author

Email address: krassivl@fmi.uni-sofia.bg (Krassimira Vlachkova)

curves for coincidence. They represented the curves into irreducible form¹, tested them for coincidence and determined their shared domain in the case where coincidence occurs. In (Berry and Patterson, 1997) a general approach and result for comparing rational Bézier curves based on their control polygons was proposed. Then, in a series of papers this approach for testing Bézier curves for coincidence has been developed and discussed, see (W.-K. Wang et al., 2011; Sánchez-Reyes, 2011; Chen et al., 2013; Sánchez-Reyes, 2014; Chen et al., 2016; Sánchez-Reyes, 2015a; Chen and Ma, 2015).

The use of control polygons when comparing Bézier curves is preferable over their reparametrization as proposed in (Pekerman et al., 2005) due to stability and numerical issues, see (Sánchez-Reyes, 2015b; Farouki, 2012). In addition, the transformations involved become ill-conditioned for high degrees. In (Vlachkova, 2017) an algorithm for testing Bézier curves for coincidence based on control polygons and subdivision was presented, analysed and experimentally tested.

Pekerman et al. (2005) suggested in the concluding remarks that their approach for testing polynomial curves for coincidence can be extended to TPB surfaces. No further detail was given. Vlachkova (2017) proposed an algorithm for testing TPB surfaces as a generalization of the algorithm for curves presented in the same paper. The algorithm is based on comparing the control polygons of the two surfaces. TPB surface $S(u, v)$, $0 \leq u \leq 1$, $0 \leq v \leq 1$, is *irreducible* if all comprising Bézier curves in u and v directions are irreducible. Two irreducible TPB surfaces can have coincident part either (i) if they are of same degree (n, m) , $n, m \in \mathbb{N}$; or (ii) if one is of degree (n, m) and the other is of degree either $(n, n + m)$, or $(n + m, m)$, or $(n + m, n + m)$. The algorithm in (Vlachkova, 2017) works only for TPB surfaces of same degree (n, m) with overlapping boundary curves, see Fig. 2a. All other cases remained open.

Here we apply a different approach and present a complete solution to the problem of comparing TPB surfaces for coincidence. First, we show that all cases where two irreducible TPB surfaces have coincidence can be reduced to two main cases: (i) the surfaces are of same degree; (ii) the surfaces are of degree (n, m) and $(n + m, n + m)$, respectively. Then, we reduce the problem to solving a nonlinear system of equations of degree $n + m$. The number of the unknowns is four and eight in cases (i) and (ii), respectively. Finally, we propose a method for solving these systems. Based on their solutions, we decide whether the input surfaces are different or partially/entirely coincident. In the latter case we determine the control points of the coincident part using the blossoming principle. We also derive sufficient geometric criteria for checking whether the surfaces are different.

We have implemented our approach using Mathematica package. We reformulate the arising nonlinear systems of high degree so that Mathematica finds correctly their solutions. The experimental results are presented, analysed and visualized.

The paper is organized as follows. In Section 2 we formulate the problem and consecutively resolve cases (i) and (ii) in Subsection 2.1 and Subsection 2.2, respectively. In Section 3 we discuss the implementation and present our experimental results. Summary and conclusions are presented in Section 4.

¹Different representations of a polynomial curve may occur if it has been degree elevated and/or reparameterized by a composition with a polynomial. A curve is *irreducible* if it is not a result of a polynomial composition and has not been degree elevated. Checking curves for irreducibility is well understood, see e. g. (Barton and Zippel, 1985; Kozen and Landau, 1989; von zur Gathen, 1990). In our experiments we use the built-in functions in Mathematica package.

2. Coincidence of TPB surfaces

Tensor product Bézier surface $S(u, v)$ of degree (n, m) for $n, m \in \mathbb{N}$, and control points $\mathbf{p}_{ij} \in \mathbb{R}^3$, $i = 0, \dots, n$, $j = 0, \dots, m$ is defined by

$$S(u, v) = \sum_{i=0}^n \sum_{j=0}^m \mathbf{p}_{ij} B_i^n(u) B_j^m(v), \quad (1)$$

where $0 \leq u \leq 1$, $0 \leq v \leq 1$, and $B_i^n(u) := \binom{n}{i} u^i (1-u)^{n-i}$ are the Bernstein polynomials. Hereafter we assume that the binomial coefficients $\binom{n}{i} = 0$ if $i < 0$ or $i > n$.

We denote by $L(S)$ the locus of points $(u, v, S(u, v))$ in \mathbb{R}^3 for $0 \leq u \leq 1$, $0 \leq v \leq 1$.

Definition 1. TPB surfaces S_1 and S_2 have *coincidence* if there exists TPB surface S such that $L(S_k) \subseteq L(S)$, $k = 1, 2$. We distinguish the following three cases.

- (i) if $L(S_1) \equiv L(S_2)$ then S_1 and S_2 are *coincident*,
- (ii) if $L(S_1) \cap L(S_2) \neq \emptyset$ then S_1 and S_2 have *coincident part*,
- (iii) if $L(S_1) \cap L(S_2) = \emptyset$ then S_1 and S_2 are *disjoint*.

S_1 and S_2 are different if they do not have coincidence.

Recall that a Bézier curve is irreducible if it is not a result of a polynomial composition and has not been degree elevated.

Definition 2. TPB surface $S(u, v)$ is *irreducible* if the Bézier curves $C_j^u(u)$, $j = 0, \dots, m$ with control points $\{\mathbf{p}_{ij}\}_{i=0}^n$, and $C_i^v(v)$, $i = 0, \dots, n$ with control points $\{\mathbf{p}_{ij}\}_{j=0}^m$ are irreducible.

For TPB surface S with control points \mathbf{p}_{ij} we denote by $\boldsymbol{\rho}$, $\boldsymbol{\rho}^{1,0}$, and $\boldsymbol{\rho}^{0,1}$ the finite differences at point \mathbf{p}_{00} of order (n, m) , $(n-1, m)$, and $(n, m-1)$ respectively, defined by (see (Farin, 2002, pp. 66, 256))

$$\begin{aligned} \boldsymbol{\rho} &:= \Delta^{n,m} \mathbf{p}_{00} = \sum_{i=0}^n \sum_{j=0}^m (-1)^{n+m-i-j} \binom{n}{i} \binom{m}{j} \mathbf{p}_{ij}, \\ \boldsymbol{\rho}^{1,0} &:= \Delta^{n-1,m} \mathbf{p}_{00} = \frac{1}{n} \sum_{i=0}^n \sum_{j=0}^m (-1)^{n+m-i-j} \binom{n}{i} \binom{m}{j} (i-n) \mathbf{p}_{ij}, \\ \boldsymbol{\rho}^{0,1} &:= \Delta^{n,m-1} \mathbf{p}_{00} = \frac{1}{m} \sum_{i=0}^n \sum_{j=0}^m (-1)^{n+m-i-j} \binom{n}{i} \binom{m}{j} (j-m) \mathbf{p}_{ij}. \end{aligned} \quad (2)$$

Remark 1. If S is irreducible then $\boldsymbol{\rho}$ is non-collinear to both $\boldsymbol{\rho}^{1,0}$ and $\boldsymbol{\rho}^{0,1}$.

It is known that two irreducible TPB surfaces S_1 and S_2 of degrees (n, m) and (n_1, m_1) , respectively, may have coincidence only if (n_1, m_1) equals to one of the following: (n, m) , $(n, n+m)$, $(n+m, m)$, $(n+m, n+m)$, see (Farin, 2002, pp. 253). We consider first the cases (i) $(n_1, m_1) = (n, m)$, $n, m \in \mathbb{N}$ and (ii) $(n_1, m_1) = (n+m, n+m)$. Then we show that the other two cases reduce to (ii).

The next proposition is shown in (Berry and Patterson, 1997) and (Sánchez-Reyes, 2011) for curves. It can be easily extended to case (i) and to case (ii) (see Theorem 1 in (Yang and Zeng, 2008)) as follows.

Proposition 1. (i) The irreducible TPB surfaces $S_1(u, v)$ and $S_2(s, t)$ of same degree (n, m) have coincidence if and only if there exists affine transformation

$$\varphi : \begin{cases} u(s) = (1-s)a + sb, \\ v(t) = (1-t)c + td, \end{cases} \quad (3)$$

such that $S_2(s, t) = S_1(u(s), v(t))$ for $0 \leq s \leq 1, 0 \leq t \leq 1$, see Fig. 1(i).

(ii) The irreducible TPB surfaces $S_1(u, v)$ and $S_2(s, t)$ of degrees (n, m) and $(n+m, n+m)$, respectively, have coincidence if and only if there exists bilinear transformation

$$\psi : \begin{cases} u(s, t) = (1-s)(1-t)a_1 + (1-s)td_1 + stc_1 + s(1-t)b_1, \\ v(s, t) = (1-s)(1-t)a_2 + (1-s)td_2 + stc_2 + s(1-t)b_2 \end{cases} \quad (4)$$

such that $S_2(s, t) = S_1(u(s, t), v(s, t))$ for $0 \leq s \leq 1, 0 \leq t \leq 1$, see Fig. 1(ii).

2.1. Irreducible TPB surfaces of same degree (n, m)

Let $S_1(u, v)$ and $S_2(s, t)$ be irreducible TPB surfaces of same degree (n, m) defined as in (1) with control points \mathbf{p}_{ij}^1 and \mathbf{p}_{ij}^2 , respectively. Let the corresponding vectors $\rho_k, \rho_k^{1,0}$, and $\rho_k^{0,1}$, $k = 1, 2$, be defined by (2). We assume that S_1 and S_2 have different control polygons. In the next lemma we derive necessary geometric conditions for S_1 and S_2 to have coincidence.

Lemma 1. *If the irreducible TPB surfaces S_1 and S_2 of degree (n, m) have coincidence then the following statements hold.*

- (i) ρ_1 and ρ_2 are collinear;
- (ii) $\rho_1, \rho_2, \rho_1^{1,0}$, and $\rho_2^{1,0}$ are coplanar;
- (iii) $\rho_1, \rho_2, \rho_1^{0,1}$, and $\rho_2^{0,1}$ are coplanar.

Proof. Assume that S_1 and S_2 have coincidence. According to statement (i) of Proposition 1 there are four numbers $a, b, c, d \in \mathbb{R}$ such that the domain $\{(u, v) : a \leq u \leq b, c \leq v \leq d\}$ is an image of the domain $\{(s, t) : 0 \leq s \leq 1, 0 \leq t \leq 1\}$ under the affine transformation (3) (see Fig. 1(i)) and $S_2(s, t) = S_1(u(s), v(t))$.




Figure 1: (i) The domain $\{(u, v) : a \leq u \leq b, c \leq v \leq d\}$ is an image of the domain $\{(s, t) : 0 \leq s \leq 1, 0 \leq t \leq 1\}$ under the affine transformation $u = u(s) = (1-s)a + sb, v = v(t) = (1-t)c + td$. (ii) The convex quadrilateral $ABCD$ is an image of the domain $\{(s, t) : 0 \leq s \leq 1, 0 \leq t \leq 1\}$ under a bilinear transformation.

We have

$$\sum_{i=0}^n \sum_{j=0}^m \mathbf{p}_{ij}^2 B_i^n(s) B_j^m(t) = \sum_{i=0}^n \sum_{j=0}^m \mathbf{p}_{ij}^1 B_i^n((1-s)a + sb) B_j^m((1-t)c + td), \quad 0 \leq s \leq 1, 0 \leq t \leq 1. \quad (5)$$

We take $\frac{\partial^{n+m}}{\partial s^n \partial t^m}$ derivative in (5) and obtain

$$n!m!\rho_2 = n!m!(b-a)^n(d-c)^m\rho_1.$$

Hence, vectors ρ_1 and ρ_2 are collinear with $\rho_2 = \kappa\rho_1$ where we denoted

$$\kappa = (b-a)^n(d-c)^m. \quad (6)$$

This proves statement (i).

Since ρ_1 and ρ_2 are the coefficients of $u^n v^m$ and $s^n t^m$ in $S_1(u, v)$ and $S_2(s, t)$, respectively, and S_1 and S_2 are irreducible then $\rho_1 \neq \mathbf{0}$ and $\rho_2 \neq \mathbf{0}$. So the number κ is determined as the ratio of any two corresponding nonzero coordinates of ρ_2 and ρ_1 .

Next, we take $\frac{\partial^{n+m-1}}{\partial s^{n-1} \partial t^m}$ derivative in (5) and obtain

$$\begin{aligned} m!(n-1)! \sum_{i=0}^n \sum_{j=0}^m (-1)^{n+m-i-j} \binom{n}{i} \binom{m}{j} (ns - n + i) \mathbf{p}_{ij}^2 = \\ m!(n-1)!(b-a)^{n-1}(d-c)^m \sum_{i=0}^n \sum_{j=0}^m (-1)^{n+m-i-j} \binom{n}{i} \binom{m}{j} (n(s(b-a) + a) - n + i) \mathbf{p}_{ij}^1, \end{aligned}$$

$$0 \leq s \leq 1,$$

which for $s = 0$ implies

$$\rho_2^{1,0} = (b-a)^{n-1}(d-c)^m(a\rho_1 + \rho_1^{1,0}). \quad (7)$$

Hence, vectors ρ_1 , $\rho_1^{1,0}$, and $\rho_2^{1,0}$ are coplanar and (ii) follows from (i).

Similarly, we take $\frac{\partial^{n+m-1}}{\partial s^n \partial t^{m-1}}$ derivative in (5) and for $s = 0$ obtain

$$\rho_2^{0,1} = (b-a)^n(d-c)^{m-1}(c\rho_1 + \rho_1^{0,1}). \quad (8)$$

Therefore vectors ρ_1 , $\rho_1^{0,1}$, and $\rho_2^{0,1}$ are coplanar and (iii) follows from (i). \square

Remark 2. Surface S_2 can be considered as obtained from surface S_1 by subdivision with respect to u at a and b , and with respect to v at c and d .

Next we obtain necessary and sufficient conditions for surfaces S_1 and S_2 to have coincidence. In the proof of Lemma 1 we have shown that if S_1 and S_2 have coincidence then there exist numbers a, b, c, d defining affine transformation (3) and satisfying the system

$$\begin{cases} \rho_2^{1,0} = (b-a)^{n-1}(d-c)^m(a\rho_1 + \rho_1^{1,0}), \\ \rho_2^{0,1} = (b-a)^n(d-c)^{m-1}(c\rho_1 + \rho_1^{0,1}), \\ \kappa = (b-a)^n(d-c)^m, \end{cases} \quad (9)$$

where κ is determined from $\rho_2 = \kappa\rho_1$.

Lemma 2. *If system (9) is consistent then it has a unique solution.*

Proof. We multiply the first vector equation in (9) by $b - a$ and the second one by $d - c$, use $\rho_2 = \kappa\rho_1$, and obtain

$$b\rho_2^{1,0} - a(\rho_2 + \rho_2^{1,0}) = \kappa\rho_1^{1,0}, \quad (10)$$

$$d\rho_2^{0,1} - c(\rho_2 + \rho_2^{0,1}) = \kappa\rho_1^{0,1}. \quad (11)$$

Hence, (9) is equivalent to the following system

$$\begin{cases} b\rho_2^{1,0} - a(\rho_2 + \rho_2^{1,0}) = \kappa\rho_1^{1,0}, \\ d\rho_2^{0,1} - c(\rho_2 + \rho_2^{0,1}) = \kappa\rho_1^{0,1}, \\ \kappa = (b - a)^n(d - c)^m, \end{cases} \quad (12)$$

Each of the two vector equations in (12) is equivalent to a linear system of three equations of the unknowns a, b , and c, d , respectively, for each of the three vector coordinates. By Remark 1 the ranks of the matrices of these systems are greater than one. Hence the systems have either unique, or no solution. Therefore, if system (12) is consistent then it has a unique solution. \square

The following theorem holds.

Theorem 1. *The irreducible TPB surfaces S_1 and S_2 of degree (n, m) have coincidence if and only if system (12) is consistent and the control polygon of the surface corresponding to the unique solution to (12) coincide with the control polygon of S_2 (up to eight different enumerations of the control polygons).*

Proof. \Rightarrow Let S_1 and S_2 have coincidence. Then according to statement (i) of Proposition 1 there exist four numbers $a, b, c, d \in \mathbb{R}$ defining transformation φ which are a solution to system (9) and $S_2(s, t) = S_1(\varphi)$. According to Lemma 2, the equivalent to (9) system (12) has a unique solution (a, b, c, d) . The control polygon of the surface corresponding to this solution coincides with the control polygon of S_2 .

\Leftarrow Let system (12) be consistent. Then, according to Lemma 2, it has a unique solution. Since the control polygon of the surface S corresponding to this solution coincide with the control polygon of S_2 (up to eight different enumerations of the control polygons) then S and S_2 coincide according to Theorem 1 in (Vlachkova, 2017). \square

We continue by providing an efficient approach for testing S_1 and S_2 for coincidence. First, we consider the two linear systems (10) and (11) which have either unique or no solution. Clearly if any of them has no solution then by Proposition 1 no transformation φ exists and S_1 and S_2 are different. If both systems have unique solutions, say (a^*, b^*) , (c^*, d^*) , then we need to check if they satisfy $(b^* - a^*)^n(d^* - c^*)^m = \kappa$. If they do not, then system (12) is inconsistent and S_1 and S_2 are different. Otherwise, following Theorem 1, we need to compute the control polygon of surface S^* corresponding to (a^*, b^*, c^*, d^*) and to check if it coincides with the control polygon of S_2 . If these polygons coincide (up to eight different enumerations of the control points) then S_1 and S_2 have coincidence, otherwise they are different. We compute the control points of S^* using the blossoming principle, see (Goldman, 2003). In (Goldman, 2003, p. 339) and (Yang and Zeng, 2008) it is pointed out that for any polynomial surface patch $S(u, v) = \sum_{i=0}^n \sum_{j=0}^m \mathbf{c}_{ij} u^i v^j$ of degree (n, m) defined for $a \leq u \leq b$ and $c \leq v \leq d$, the Bézier control points $\mathbf{p}_{v\mu}$, $v = 0, \dots, n$, $\mu = 0, \dots, m$, of this surface patch are

$$\mathbf{p}_{v\mu} = b^{\square}(\underbrace{a, \dots, a}_v, \underbrace{b, \dots, b}_{n-v}, \underbrace{c, \dots, c}_\mu, \underbrace{d, \dots, d}_{m-\mu}), \quad (13)$$

where $b^{\square}(u_1, \dots, u_n, v_1, \dots, v_m) = \sum_{i=0}^n \sum_{j=0}^m \mathbf{c}_{ij} b_{ij}^{\square}(u_1, \dots, u_n, v_1, \dots, v_m)$ is the blossom of $S(u, v)$, and

$$b_{ij}^{\square}(u_1, \dots, u_n, v_1, \dots, v_m) = \sum_{\{\alpha_1, \dots, \alpha_i\} \subseteq \{1, \dots, n\}} \frac{u_{\alpha_1} \dots u_{\alpha_i}}{\binom{n}{i}} \sum_{\{\beta_1, \dots, \beta_j\} \subseteq \{1, \dots, m\}} \frac{v_{\beta_1} \dots v_{\beta_j}}{\binom{m}{j}} \quad (14)$$

is the blossom of the monomial $u^i v^j$. In the case where either $i = 0$, or $j = 0$, the corresponding sum in (14) equals 1, e. g. $b_{00}^\square = 1$. In the next corollary we present (13) and (14) in an equivalent closed form that is more suitable for computations.

Corollary 1. Bézier control points $\mathbf{p}_{v\mu}$, $v = 0, \dots, n$, $\mu = 0, \dots, m$, defined by (13) are

$$\mathbf{p}_{v\mu} = \sum_{i=0}^n \sum_{j=0}^m \frac{\mathbf{c}_{ij}}{\binom{n}{i} \binom{m}{j}} \sum_{k=\max(0, i+v-n)}^{\min(i, v)} \sum_{r=\max(0, j+\mu-m)}^{\min(j, \mu)} \binom{v}{k} \binom{n-v}{i-k} \binom{\mu}{r} \binom{m-\mu}{j-r} a^k b^{i-k} c^r d^{j-r}. \quad (15)$$

We outline our procedure for testing S_1 and S_2 for coincidence in algorithmic form below.

Algorithm 1 Testing two irreducible TPB surfaces of degree (n, m) for coincidence

Input: Irreducible TPB surfaces S_1 and S_2 of degree (n, m) given by their control polygons

Output: (i) S_1 and S_2 are different;
(ii) S_1 and S_2 are disjoint;
(iii) S_1 and S_2 have coincident part. Report its control points.

Step 1. Compute vectors ρ_i , $\rho_i^{1,0}$, and $\rho_i^{0,1}$, $i = 1, 2$.

Step 2. Check the conditions of Lemma 1.

2.1. **If** ρ_1, ρ_2 are non-collinear

then return (i);

else compute κ such that $\rho_2 = \kappa\rho_1$.

2.2. **If** either $\rho_1, \rho_2, \rho_1^{1,0}, \rho_2^{1,0}$, or $\rho_1, \rho_2, \rho_1^{0,1}, \rho_2^{0,1}$ are non-coplanar

then return (i);

else system (12) has either unique, or no solution.

Step 3. Solve linear systems (10) and (11).

If any of them is inconsistent

then return (i);

else denote their unique solutions by (a^*, b^*) and (c^*, d^*) .

Step 4. **If** $(b^* - a^*)^n (d^* - c^*)^m \neq \kappa$

then return (i);

else system (9) is consistent with unique solution (a^*, b^*, c^*, d^*) .

Step 5. Compute the control polygon of the transformed TPB surface

$S^*(s, t) = S_1(u(\varphi(s, t), v(\varphi(s, t)))$ using (15) and compare it to the control polygon of S_2 .

If they coincide (up to eight different enumerations)

then S_1 and S_2 have coincidence;

else return (i).

Step 6. Compute the shared domain G of S^* and S .

If $D = \emptyset$

then return (ii);

else compute the control points of the coincident part using (15) and return (iii).

2.2. Irreducible Bézier surfaces of degrees (n, m) and $(n + m, n + m)$

Let $S_1(u, v) = \sum_{i=0}^n \sum_{j=0}^m \mathbf{p}_{ij}^1 B_i^n(u) B_j^m(v)$ defined for $0 \leq u \leq 1, 0 \leq v \leq 1$, and $S_2(s, t) = \sum_{i=0}^{n+m} \sum_{j=0}^{n+m} \mathbf{p}_{ij}^2 B_i^{n+m}(s) B_j^{n+m}(t)$ defined for $0 \leq s \leq 1, 0 \leq t \leq 1$, be irreducible TPB surfaces. For the four boundary curves of S_2 we denote by δ_k and

$\delta_k^1, k = 1, \dots, 4$ the following finite differences

$$\begin{aligned}\delta_1 &:= \Delta^{n+m,0} \mathbf{p}_{00}^2 = \sum_{i=0}^{n+m} (-1)^{n+m-i} \binom{n+m}{i} \mathbf{p}_{i0}^2, \\ \delta_2 &:= \Delta^{0,n+m} \mathbf{p}_{00}^2 = \sum_{i=0}^{n+m} (-1)^{n+m-i} \binom{n+m}{i} \mathbf{p}_{0i}^2, \\ \delta_3 &:= \Delta^{n+m,0} \mathbf{p}_{0,n+m}^2 = \sum_{i=0}^{n+m} (-1)^{n+m-i} \binom{n+m}{i} \mathbf{p}_{i,n+m}^2, \\ \delta_4 &:= \Delta^{0,n+m} \mathbf{p}_{n+m,0}^2 = \sum_{i=0}^{n+m} (-1)^{n+m-i} \binom{n+m}{i} \mathbf{p}_{n+m,i}^2, \\ \delta_1^1 &:= (n+m) \Delta^{n+m-1,0} \mathbf{p}_{00}^2 = \sum_{i=0}^{n+m} (-1)^{n+m-i} \binom{n+m}{i} (i-n-m) \mathbf{p}_{i0}^2, \\ \delta_2^1 &:= (n+m) \Delta^{0,n+m-1} \mathbf{p}_{00}^2 = \sum_{i=0}^{n+m} (-1)^{n+m-i} \binom{n+m}{i} (i-n-m) \mathbf{p}_{0i}^2, \\ \delta_3^1 &:= (n+m) \Delta^{n+m-1,0} \mathbf{p}_{0,n+m}^2 = \sum_{i=0}^{n+m} (-1)^{n+m-i} \binom{n+m}{i} (i-n-m) \mathbf{p}_{i,n+m}^2, \\ \delta_4^1 &:= (n+m) \Delta^{0,n+m-1} \mathbf{p}_{n+m,0}^2 = \sum_{i=0}^{n+m} (-1)^{n+m-i} \binom{n+m}{i} (i-n-m) \mathbf{p}_{n+m,i}^2.\end{aligned}$$

The finite differences $\rho, \rho^{1,0}$, and $\rho^{0,1}$ for S_1 are defined by (2).

In the next lemma we derive necessary geometric conditions for S_1 and S_2 to have coincidence.

Lemma 3. *If the irreducible TPB surfaces S_1 and S_2 of degrees (n, m) and $(n+m, n+m)$, respectively, have coincidence then the five vectors ρ , and $\delta_i, i = 1, \dots, 4$, are collinear.*

Proof. Assume that S_1 and S_2 have coincidence. Since S_1 is of degree (n, m) and S_2 is of degree $(n+m, n+m)$ then, according to statement (ii) of Proposition 1, there exist convex quadrilateral $ABCD$ with vertices $A(a_1, a_2), B(b_1, b_2), C(c_1, c_2), D(d_1, d_2)$ which is an image of the domain $\{(s, t) : 0 \leq s \leq 1, 0 \leq t \leq 1\}$ under the bilinear transformation (4) (see Fig. 1(ii)) and $S_2(s, t) = S_1(u(s, t), v(s, t))$. Hence, for $0 \leq s \leq 1, 0 \leq t \leq 1$ we have

$$\sum_{i=0}^n \sum_{j=0}^m \mathbf{p}_{ij}^1 B_i^n(u) B_j^m(v) = \sum_{i=0}^{n+m} \sum_{j=0}^{n+m} \mathbf{p}_{ij}^2 B_i^{n+m}(s) B_j^{n+m}(t), \quad (16)$$

where u and v are defined by (4).

The image of the boundary segment $\{(s, t) : 0 \leq s \leq 1, t = 0\}$ under the bilinear transformation (4) is the segment AB and we have $u = (1-s)a_1 + sb_1, v = (1-s)a_2 + sb_2$. Hence, from (16) it follows

$$\sum_{i=0}^n \sum_{j=0}^m \mathbf{p}_{ij}^1 B_i^n((1-s)a_1 + sb_1) B_j^m((1-s)a_2 + sb_2) = \sum_{i=0}^{n+m} \mathbf{p}_{i0}^2 B_i^{n+m}(s), \quad 0 \leq s \leq 1. \quad (17)$$

After differentiation of (17) $n+m$ times we obtain

$$(b_1 - a_1)^n (b_2 - a_2)^m \rho = \delta_1. \quad (18)$$

Hence, vectors ρ and δ_1 are collinear with $\kappa_1\rho = \delta_1$ where we denoted

$$\kappa_1 = (b_1 - a_1)^n(b_2 - a_2)^m. \quad (19)$$

Since ρ and δ_1 are the coefficients of $u^n v^m$ in $S_1(u, v)$ and s^{n+m} in $S_2(s, 0)$, respectively, and S_1 and S_2 are irreducible then $\rho \neq \mathbf{0}$ and $\delta_1 \neq \mathbf{0}$. So the number κ_1 is determined as the ratio of any two corresponding nonzero coordinates of δ_1 and ρ .

For the remaining three boundary segments we obtain, analogously to (18),

$$\begin{aligned} (d_1 - a_1)^n(d_2 - a_2)^m \rho &= \delta_2, \\ (c_1 - d_1)^n(c_2 - d_2)^m \rho &= \delta_3, \\ (c_1 - b_1)^n(c_2 - b_2)^m \rho &= \delta_4. \end{aligned} \quad (20)$$

Hence, vectors ρ , δ_2 , δ_3 , and δ_4 are collinear with $\kappa_i\rho = \delta_i$, $i = 2, 3, 4$, where we denoted

$$\begin{aligned} \kappa_2 &= (d_1 - a_1)^n(d_2 - a_2)^m, \\ \kappa_3 &= (c_1 - d_1)^n(c_2 - d_2)^m, \\ \kappa_4 &= (c_1 - b_1)^n(c_2 - b_2)^m. \end{aligned} \quad (21)$$

Similarly to κ_1 , the numbers κ_i for $i = 2, 3, 4$ are determined as the ratio of any two corresponding nonzero coordinates of δ_i and ρ , respectively. \square

Next we obtain a necessary and sufficient conditions for surfaces S_1 and S_2 to have coincidence. We differentiate (17) $n + m - 1$ times and obtain for $s = 0$

$$(b_1 - a_1)^{n-1}(b_2 - a_2)^{m-1}(n\rho^{1,0}(b_2 - a_2) + m\rho^{0,1}(b_1 - a_1) + n\rho a_1(b_2 - a_2) + m\rho a_2(b_1 - a_1)) = \delta_1^1.$$

Similarly, for the remaining three boundary segments we obtain

$$\begin{aligned} (d_1 - a_1)^{n-1}(d_2 - a_2)^{m-1}(n\rho^{1,0}(d_2 - a_2) + m\rho^{0,1}(d_1 - a_1) + n\rho a_1(d_2 - a_2) + m\rho a_2(d_1 - a_1)) &= \delta_2^1, \\ (c_1 - d_1)^{n-1}(c_2 - d_2)^{m-1}(n\rho^{1,0}(c_2 - d_2) + m\rho^{0,1}(c_1 - d_1) + n\rho d_1(c_2 - d_2) + m\rho d_2(c_1 - d_1)) &= \delta_3^1, \\ (c_1 - b_1)^{n-1}(c_2 - b_2)^{m-1}(n\rho^{1,0}(c_2 - b_2) + m\rho^{0,1}(c_1 - b_1) + n\rho b_1(c_2 - b_2) + m\rho b_2(c_1 - b_1)) &= \delta_4^1. \end{aligned}$$

Therefore, if S_1 and S_2 have coincidence then the eight numbers a_i, b_i, c_i, d_i , $i = 1, 2$, defining bilinear transformation (4) satisfy the system

$$\left\{ \begin{aligned} (b_1 - a_1)^{n-1}(b_2 - a_2)^{m-1}(n\rho^{1,0}(b_2 - a_2) + m\rho^{0,1}(b_1 - a_1) + n\rho a_1(b_2 - a_2) + m\rho a_2(b_1 - a_1)) &= \delta_1^1, \\ (d_1 - a_1)^{n-1}(d_2 - a_2)^{m-1}(n\rho^{1,0}(d_2 - a_2) + m\rho^{0,1}(d_1 - a_1) + n\rho a_1(d_2 - a_2) + m\rho a_2(d_1 - a_1)) &= \delta_2^1, \\ (c_1 - d_1)^{n-1}(c_2 - d_2)^{m-1}(n\rho^{1,0}(c_2 - d_2) + m\rho^{0,1}(c_1 - d_1) + n\rho d_1(c_2 - d_2) + m\rho d_2(c_1 - d_1)) &= \delta_3^1, \\ (c_1 - b_1)^{n-1}(c_2 - b_2)^{m-1}(n\rho^{1,0}(c_2 - b_2) + m\rho^{0,1}(c_1 - b_1) + n\rho b_1(c_2 - b_2) + m\rho b_2(c_1 - b_1)) &= \delta_4^1, \\ \kappa_1 &= (b_1 - a_1)^n(b_2 - a_2)^m, \\ \kappa_2 &= (d_1 - a_1)^n(d_2 - a_2)^m, \\ \kappa_3 &= (c_1 - d_1)^n(c_2 - d_2)^m, \\ \kappa_4 &= (c_1 - b_1)^n(c_2 - b_2)^m, \end{aligned} \right. \quad (22)$$

where κ_i is determined from $\kappa_i \rho = \delta_i$, $i = 1, \dots, 4$.

Straightforward application of Mathematica packages and build-in functions doesn't yield solutions to system (22) efficiently. Hence, it is important to develop a method to simplify and solve it.

Lemma 4. *If system (22) is consistent and $\text{rank}(\rho, \rho^{1,0}, \rho^{0,1})=3$ then it has a unique solution. If $\text{rank}(\rho, \rho^{1,0}, \rho^{0,1})=2$ then it has at most two solutions.*

Proof. We consider the first equation of system (22), denote $x := b_1 - a_1$, $y := b_2 - a_2$, and obtain

$$x^{n-1}y^{m-1}(n\rho^{1,0}y + m\rho^{0,1}x + n\rho a_1y + m\rho a_2x) = \delta_1^1. \quad (23)$$

Vector equation (23) is equivalent to the following system

$$\begin{cases} \kappa_1(n\rho_1^{1,0}y + m\rho_1^{0,1}x + n\rho_1 a_1y + m\rho_1 a_2x) = \delta_1 xy, \\ \kappa_1(n\rho_2^{1,0}y + m\rho_2^{0,1}x + n\rho_2 a_1y + m\rho_2 a_2x) = \delta_2 xy, \\ \kappa_1(n\rho_3^{1,0}y + m\rho_3^{0,1}x + n\rho_3 a_1y + m\rho_3 a_2x) = \delta_3 xy, \end{cases} \quad (24)$$

where $\rho^{1,0} = (\rho_1^{1,0}, \rho_2^{1,0}, \rho_3^{1,0})$, $\rho^{0,1} = (\rho_1^{0,1}, \rho_2^{0,1}, \rho_3^{0,1})$, $\rho = (\rho_1, \rho_2, \rho_3)$, and $\delta_1^1 = (\delta_1, \delta_2, \delta_3)$.

To solve (24) we consider two cases according to the rank of the matrix M , where

$$M = \begin{pmatrix} n\rho_1^{1,0} & m\rho_1^{0,1} & n\rho_1 & m\rho_1 \\ n\rho_2^{1,0} & m\rho_2^{0,1} & n\rho_2 & m\rho_2 \\ n\rho_3^{1,0} & m\rho_3^{0,1} & n\rho_3 & m\rho_3 \end{pmatrix}. \quad (25)$$

By Remark 1 the rank of M is greater than 1. Further on, $\alpha, \beta, C, \alpha_i, \beta_i, C_i$, $i = 1, 2$ denote real constants that depend on the input data only, more precisely on $\rho^{1,0}$, $\rho^{0,1}$, ρ , and δ_1^1 .

Case 1. $\text{rank}(M)=3$

Since ρ is nonzero vector then some of its coordinates, say ρ_1 , is nonzero. We eliminate a_1y and a_2x from the last two equations of (24) by multiplying the first equation by $-\rho_2/\rho_1$ and $-\rho_3/\rho_1$ consecutively and adding it to the second and third equations, respectively. We obtain a system of the following type

$$\begin{cases} \alpha_1 x + \beta_1 y = C_1 xy, \\ \alpha_2 x + \beta_2 y = C_2 xy, \end{cases} \quad (26)$$

which has a unique solution (x, y) .

We solve in an analogous way the remaining three vector equations of (22) with respect to the unknowns $d_1 - a_1$ and $d_2 - a_2$; $c_1 - d_1$ and $c_2 - d_2$; $c_1 - b_1$ and $c_2 - b_2$, respectively. Note that the corresponding three equivalent systems have same matrix M as system (24) and differ by their right sides only. Hence, each of them has also a unique solution which can be found straightforwardly. Therefore, if system (22) is consistent then it has a unique solution.

Case 2. $\text{rank}(M)=2$

In this case, (24) has two linearly independent equations. Similar to Case 1, since the coefficients of a_1y and a_2x are in ratio $n : m$, then by multiplying one of these equations by a suitable constant and adding it to the other equation we exclude a_1y and a_2x and obtain one equation of the following type

$$\alpha x + \beta y = C xy. \quad (27)$$

So we have to solve the following system

$$\begin{cases} x^n y^m = \kappa_1, \\ \alpha x + \beta y = Cxy. \end{cases} \quad (28)$$

Claim 1. System (28) has at most two solutions in real numbers.

Proof. Omitted.

In the general case where $\alpha \neq 0$ and $\beta \neq 0$ from (27) we have $y = \alpha x / (Cx - \beta)$. Hence, system (28) reduces to the following polynomial equation of degree $n + m$

$$x^n (\alpha x)^m = \kappa_1 (Cx - \beta)^m. \quad (29)$$

We solve (29) using Mathematica and find all its real solutions. We note that if $n + m$ is odd then the solution is unique, otherwise (29) may have two solutions.

So far we have found all admissible values for the unknown $x = b_1 - a_1$ and $y = b_2 - a_2$. We solve in an analogous way the remaining three vector equations of (22) with respect to the unknowns $d_1 - a_1$ and $d_2 - a_2$; $c_1 - d_1$ and $c_2 - d_2$; $c_1 - b_1$ and $c_2 - b_2$, respectively. Recall that the four systems have same matrix M and differ by their right sides only.

Further, we select all combinations of quadruplets (x_1, x_2, x_3, x_4) and (y_1, y_2, y_3, y_4) such that (x_i, y_i) , $i = 1, \dots, 4$ are solutions to the first four equations of (22), respectively, and in addition satisfy the following conditions

$$x_1 - x_2 - x_3 + x_4 = 0, \quad y_1 - y_2 - y_3 + y_4 = 0. \quad (30)$$

Next we show how to obtain the eight unknowns a_i, b_i, c_i, d_i , $i = 1, 2$, from the selected quadruplets (x_1, x_2, x_3, x_4) and (y_1, y_2, y_3, y_4) , if any. Let (x_1, x_2, x_3, x_4) and (y_1, y_2, y_3, y_4) be a couple of the selected quadruplets. Since b_i, c_i, d_i , $i = 1, 2$, can be represented through a_1, a_2 as

$$b_1 = a_1 + x_1, b_2 = a_2 + y_1; c_1 = a_1 + x_2 + x_3, c_2 = a_2 + y_2 + y_3; d_1 = a_1 + x_1, d_2 = a_2 + y_2, \quad (31)$$

then we have to find a_1, a_2 only. We replace (x_i, y_i) , $i = 1, \dots, 4$, and the relations (31) in the four vector equations of (22) and for each of them we obtain a linear equation of a_1 and a_2 . If the system of these four linear equations is consistent, i. e. its rank is 2, and in addition the corresponding a_i, b_i, c_i, d_i , $i = 1, 2$, satisfy the last four equations of (22) then $(a_1, a_2, b_1, b_2, c_1, c_2, d_1, d_2)$ is a solution to (22). Otherwise, the selected couple of quadruplets does not produce a solution to (22). In this case, if system (22) is consistent it may have at most two solutions and we have shown how to find both of them. \square

The following theorem holds.

Theorem 2. *The irreducible TPB surfaces S_1 and S_2 of degrees (n, m) and $(n+m, n+m)$, respectively, have coincidence if and only if system (22) is consistent and the control polygon of the surface corresponding to any solution to (22) coincide with the control polygon of S_2 (up to eight different enumeration of the control points).*

Proof. \Rightarrow Let S_1 and S_2 have coincidence. Then according to statement (ii) of Proposition 1 there exist eight numbers $a_i, b_i, c_i, d_i \in \mathbb{R}$, $i = 1, 2$, defining transformation ψ which are a solution to system (22) and $S_2(s, t) = S_1(\psi)$. The control polygon of the surface corresponding to the solution $(a_1, a_2, b_1, b_2, c_1, c_2, d_1, d_2)$ coincides with the control polygon of S_2 .

\Leftarrow Let system (22) be consistent. Then, according to Lemma 4, it may have at most two solutions. Since there is a solution such that the control polygon of the surface S corresponding to this solution coincides with the control polygon of S_2 (up to eight different enumerations of the control polygons) then S and S_2 coincide according to Theorem 1 in (Vlachkova, 2017). \square

We continue by providing an efficient approach for testing S_1 and S_2 for coincidence. First, we find all real solutions to system (22) using the proposed technique in Lemma 4. Clearly, if system (22) has no real solution then by Proposition 1 no transformation ψ exists and S_1 and S_2 are different. Otherwise, let $(a_1, a_2, b_1, b_2, c_1, c_2, d_1, d_2)$ be a solution to system (22). Following Theorem 2, we need to compute the control polygon of surface S corresponding to this solution and to check if it coincides with the control polygon of S_2 . If these polygons coincide (up to eight different enumerations of the control points) then S_1 and S_2 have coincidence, otherwise they are different and we continue by checking the second solution to system (22), if any.

Next we describe how we compute the control points of the corresponding surface S defined in quadrangle $\square ABCD$ with vertices $A(a_1, b_1)$, $B(b_1, b_2)$, $C(c_1, c_2)$, and $D(d_1, d_2)$ by (13). First, we compute the shared domain G of S and S_2 . If G is the empty set then S_1 and S_2 are disjoint. Otherwise S_1 and S_2 have coincident part and we find it by using the blossoming principle. In this case, unlike the case where S_1 and S_2 are of same degree, the shared domain can be a polygon with at most eight vertices, see Fig. 6. If the number of the polygon vertices is even we represent the coincident part as a union of TPB surfaces. For example, the surface in Fig. 6c. is represented as a union of two TPB surfaces. If the number of the polygon vertices is odd we represent the coincident part as a union of TPB surfaces and a triangular Bézier (TB) surface. Similar to TPB surface, we compute the control points of the TB surface using the blossoming principle. In (Goldman, 2003, p. 331) and (Yang and Zeng, 2008) it is pointed out that for any polynomial surface path $S(u, v) = \sum_{i=0}^n \sum_{j=0}^m \mathbf{c}_{ij} u^i v^j$ of total degree $n + m$ defined in a triangle $\triangle MNP$, the Bézier control points $\mathbf{q}_{\nu\mu}$ of this surface patch are

$$\mathbf{q}_{\nu\mu} = b^\Delta(\underbrace{M, \dots, M}_\nu, \underbrace{N, \dots, N}_\mu, \underbrace{P, \dots, P}_{n+m-\nu-\mu}), \quad (32)$$

where $b^\Delta((u_1, v_1), \dots, (u_{n+m}, v_{n+m})) = \sum_{i=0}^n \sum_{j=0}^m \mathbf{c}_{ij} b_{ij}^\Delta((u_1, v_1), \dots, (u_{n+m}, v_{n+m}))$ is the blossom of $S(u, v)$, and

$$b_{ij}^\Delta((u_1, v_1), \dots, (u_{n+m}, v_{n+m})) = \sum_{\substack{\{\alpha_1, \dots, \alpha_i\} \subseteq K \\ \{\beta_1, \dots, \beta_j\} \subseteq K \setminus \{\alpha_1, \dots, \alpha_i\}}} \frac{u_{\alpha_1} \dots u_{\alpha_i} v_{\beta_1} \dots v_{\beta_j}}{\binom{n+m}{i, j}}, \quad (33)$$

$K = \{1, \dots, n + m\}$, $\binom{n+m}{i, j} = \frac{(n+m)!}{i!j!(n+m-i-j)!}$, is the blossom of the monomial $u^i v^j$. In the case where either $i = 0$, or $j = 0$ the corresponding sum in (33) equals 1. In the next corollary we present (32) in an equivalent closed form that is more suitable for evaluations of the control points.

Corollary 2. Bézier control points $\mathbf{q}_{\nu\mu}$, $\nu = 0, \dots, n$, $\mu = 0, \dots, m$, defined by (32) are

$$\mathbf{q}_{\nu\mu} = \sum_{i=0}^n \sum_{j=0}^m \frac{\mathbf{c}_{ij}}{\binom{n+m}{i, j}} b_{ij}^\Delta = \sum_{j_\beta=j_\beta}^{\bar{j}_\beta} \sum_{j_\alpha=j_\alpha}^{\bar{j}_\alpha} \sum_{i_\beta=i_\beta}^{\bar{i}_\beta} \sum_{i_\alpha=i_\alpha}^{\bar{i}_\alpha} \binom{\nu}{i_\alpha} \binom{\mu}{i_\beta} \binom{\lambda}{i_\gamma} \binom{\nu-i_\alpha}{j_\alpha} \binom{\mu-i_\beta}{j_\beta} \binom{\lambda-i_\gamma}{j_\gamma} a_1^{i_\alpha} b_1^{i_\beta} c_1^{i_\gamma} a_2^{j_\alpha} b_2^{j_\beta} c_2^{j_\gamma}, \quad (34)$$

where

$$\begin{aligned}
\lambda &= n + m - \nu - \mu, \quad i_\gamma = i - i_\alpha - i_\beta, \quad j_\gamma = j - j_\alpha - j_\beta, \\
\underline{i}_\alpha &= \max(0, i - \mu - \lambda), \quad \bar{i}_\alpha = \min(i, \nu), \\
\underline{i}_\beta &= \max(0, i - \nu - \lambda), \quad \bar{i}_\beta = \min(i - i_\alpha, \mu), \\
\underline{j}_\alpha &= \max(0, j - (\mu - i_\beta) - (\lambda - i_\gamma)), \quad \bar{j}_\alpha = \min(j, \nu - i_\alpha), \\
\underline{j}_\beta &= \max(0, j - (\nu - i_\alpha) - (\lambda - i_\gamma)), \quad \bar{j}_\beta = \min(j - j_\alpha, \mu - i_\beta).
\end{aligned}$$

We outline our procedure for testing S_1 and S_2 for coincidence in algorithmic form in Algorithm 2.

Algorithm 2 Testing two irreducible TPB surfaces of degree (n, m) and $(n + m, n + m)$ for coincidence

Input: Irreducible TPB surfaces S_1 and S_2 of degree (n, m) and $(n + m, n + m)$ given by their control polygons

Output: (i) S_1 and S_2 are different;

(ii) S_1 and S_2 are disjoint;

(iii) S_1 and S_2 have coincident part. Report its control points.

Step 1. Compute vectors $\rho, \rho^{1,0}, \rho^{0,1}, \delta_i,$ and $\delta_i^1, i = 1, \dots, 4.$

Step 2. **If** ρ and δ_i for any $i, i = 1, \dots, 4,$ are non-collinear

then return (i);

else compute κ_i such that $\kappa_i \rho = \delta_i, i = 1, \dots, 4.$

Step 3. Solve the first four vector equations of (22) according to Case 1 and Case 2 of Lemma 4 and find all admissible couples of quadruplets (x_1, x_2, x_3, x_4) and $(y_1, y_2, y_3, y_4).$

Step 4. For any admissible couple of quadruplets

if either $x_1 - x_2 - x_3 + x_4 \neq 0,$ or $y_1 - y_2 - y_3 + y_4 \neq 0$

then return (i);

else if $x_i^n y_i^m \neq k_i$ for any $i, i = 1, \dots, 4$

then return (i);

else substitute (31) in the vector equations of (22) and obtain a system of four linear equations of a_1 and $a_2.$

If this system is inconsistent

then return (i);

else $(a_1, a_2, b_1, b_2, c_1, c_2, d_1, d_2)$ is a solution to (22).

Step 5. Compute the control polygon of the transformed TPB surface $S(s, t) = S_1(\psi(s, t))$ using (15) and (34) and compare it to the control polygon of $S_2.$

If they coincide (up to eight different enumerations)

then S_1 and S_2 have coincidence;

else return (i).

Step 6. Compute the shared domain G of S and $S_2.$

If $D = \emptyset$

then return (ii);

else divide G into quadrangles and a triangle (if necessary), compute the control points of the corresponding coincident Bézier surfaces using (15) and (34), and return (iii).

Remark 3. In the case where the shared domain is not a rectangle, multiple representations of the coincident part as a union of Bézier surfaces exist.

Remark 4. The case where S_1 is of degree (n, m) and S_2 is of degree either $(n, n + m)$, or $(n + m, m)$ is analogous to the case where S_2 is of degree $(n + m, n + m)$. The only difference is that (16) becomes (e. g. for degree $(n + m, m)$)

$$\sum_{i=0}^n \sum_{j=0}^m \mathbf{p}_{ij}^1 B_i^n(u) B_j^m(v) = \sum_{i=0}^{n+m} \sum_{j=0}^m \mathbf{p}_{ij}^2 B_i^{n+m}(s) B_j^m(t).$$

Further, the arguments are the same as in the case where S_1 is of degree (n, m) and S_2 is of degree $(n + m, n + m)$.

3. Examples and results

We have implemented our method using Mathematica package. In this section we present and analyze the results from our experimental work. In our examples we consider irreducible TPB surfaces.

Example 1. This example illustrates the only case where the direct generalization of the algorithm for curves works, see (Vlachkova, 2017). More precisely, this is the case where the two TPB surfaces of same degree have overlapping boundary curves. Here we apply the new method outlined in Algorithm 1. The irreducible surfaces S_1 and S_2 are of degree $(4, 2)$. Their control points are shown in Table 1. The unique solution of system (9) is $(a^*, b^*, c^*, d^*) = (0, 1/2, 1/6, 3/4)$. Surfaces S_1 and S_2 are shown in Fig. 2a. and Fig. 2b., respectively. Their coincident part is surface S_2 . Both surfaces and their coincident part with its control polygon are shown in Fig. 3a.




Figure 2: (illustrates examples 1 and 2) Three irreducible TPB surfaces of degree $(4, 2)$ with control points shown in Table 1. **a.** Surface S_1 ; **b.** Surface S_2 ; **c.** Surface S_3 ;

Example 2. We test for coincidence the irreducible surfaces S_1 and S_3 of degree $(4, 2)$ with control points shown in Table 1. We apply Algorithm 1. The unique solution of system (9) is $(a^*, b^*, c^*, d^*) = (-1/5, 1/2, 1/6, 3/4)$. Surfaces S_1 and S_3 are shown in Fig. 2b. and Fig. 2c., respectively. Their coincident part is S_2 and it is shown with its control polygon in Fig. 2b.




Figure 3: Testing for coincidence the surfaces shown in Fig. 2. **a.** Example 1: The coincident part of S_1 and S_2 is S_2 ; **b.** Example 2: The coincident part of S_1 and S_3 is S_2 .

Table 1: The control points of irreducible TPB surfaces S_1, S_2, S_3 of degree (4, 2) from examples 1 and 2

S_1	$\mathbf{p}_{00} = (0, 0, 0)$	$\mathbf{p}_{01} = (0, \frac{7}{10}, -\frac{1}{2})$	$\mathbf{p}_{02} = (0, 2, -\frac{4}{5})$
	$\mathbf{p}_{10} = (1, 0, 1)$	$\mathbf{p}_{11} = (1, 1, 0)$	$\mathbf{p}_{12} = (1, 1, 1)$
	$\mathbf{p}_{20} = (2, 0, \frac{3}{4})$	$\mathbf{p}_{21} = (2, 1, \frac{6}{5})$	$\mathbf{p}_{22} = (2, 2, \frac{3}{4})$
	$\mathbf{p}_{30} = (3, -1, \frac{3}{20})$	$\mathbf{p}_{31} = (3, 1, \frac{2}{5})$	$\mathbf{p}_{32} = (3, 2, \frac{1}{5})$
	$\mathbf{p}_{40} = (4, 0, \frac{1}{5})$	$\mathbf{p}_{41} = (4, 2, \frac{1}{4})$	$\mathbf{p}_{42} = (4, 3, \frac{3}{4})$
S_2	$\mathbf{p}_{00} = (\frac{2}{3}, -\frac{5}{324}, \frac{6157}{12960})$	$\mathbf{p}_{01} = (\frac{2}{3}, \frac{2179}{5184}, \frac{9829}{51840})$	$\mathbf{p}_{02} = (\frac{2}{3}, \frac{1073}{1296}, \frac{9689}{103680})$
	$\mathbf{p}_{10} = (\frac{5}{4}, \frac{-23}{432}, \frac{425}{576})$	$\mathbf{p}_{11} = (\frac{5}{4}, \frac{1567}{3456}, \frac{5881}{11520})$	$\mathbf{p}_{12} = (\frac{5}{4}, \frac{1447}{1728}, \frac{6673}{13824})$
	$\mathbf{p}_{20} = (\frac{11}{6}, -\frac{1}{6}, \frac{3907}{5760})$	$\mathbf{p}_{21} = (\frac{11}{6}, \frac{107}{256}, \frac{15241}{23040})$	$\mathbf{p}_{22} = (\frac{11}{6}, \frac{515}{576}, \frac{30277}{46080})$
	$\mathbf{p}_{30} = (\frac{29}{12}, -\frac{27}{64}, \frac{1837}{3840})$	$\mathbf{p}_{31} = (\frac{29}{12}, \frac{165}{512}, \frac{2837}{5120})$	$\mathbf{p}_{32} = (\frac{29}{12}, \frac{703}{768}, \frac{17401}{30720})$
	$\mathbf{p}_{40} = (3, -\frac{27}{64}, \frac{849}{2560})$	$\mathbf{p}_{41} = (3, \frac{2287}{5120}, \frac{4253}{10240})$	$\mathbf{p}_{42} = (3, \frac{1433}{1280}, \frac{9489}{20480})$
S_3	$\mathbf{p}_{00} = (\frac{2}{3}, -\frac{11941}{32400}, \frac{475469}{648000})$	$\mathbf{p}_{01} = (\frac{2}{3}, \frac{247}{960}, \frac{29557}{129600})$	$\mathbf{p}_{02} = (\frac{2}{3}, \frac{1073}{1296}, \frac{9689}{103680})$
	$\mathbf{p}_{10} = (\frac{5}{4}, -\frac{3443}{7200}, \frac{411113}{432000})$	$\mathbf{p}_{11} = (\frac{5}{4}, \frac{1727}{5760}, \frac{11267}{21600})$	$\mathbf{p}_{12} = (\frac{5}{4}, \frac{1447}{1728}, \frac{6673}{13824})$
	$\mathbf{p}_{20} = (\frac{11}{6}, -\frac{1877}{2880}, \frac{199789}{288000})$	$\mathbf{p}_{21} = (\frac{11}{6}, \frac{2621}{11520}, \frac{849}{1280})$	$\mathbf{p}_{22} = (\frac{11}{6}, \frac{515}{576}, \frac{30277}{46080})$
	$\mathbf{p}_{30} = (\frac{29}{12}, -\frac{9997}{9600}, \frac{26091}{64000})$	$\mathbf{p}_{31} = (\frac{29}{12}, \frac{653}{7680}, \frac{659}{1200})$	$\mathbf{p}_{32} = (\frac{29}{12}, \frac{703}{768}, \frac{17401}{30720})$
	$\mathbf{p}_{40} = (3, -\frac{36737}{32000}, \frac{33149}{128000})$	$\mathbf{p}_{41} = (3, \frac{909}{5120}, \frac{10141}{25600})$	$\mathbf{p}_{42} = (3, \frac{1433}{1280}, \frac{9489}{20480})$




Figure 4: The irreducible TPB surfaces S_1 of degree (2, 3) and S_2 of degree (5, 5) from Example 3. The corresponding control points are shown in Table 2. **a.** Surface S_1 ; **b.** Surface S_2 ; **c.** The coincident part of S_1 and S_2 is S_2 .

Table 2: The control points of irreducible TPB surfaces of degree (2, 3) and (5, 5) from Example 3

S_1	$\mathbf{p}_{00} = (0, 0, 0)$	$\mathbf{p}_{01} = (\frac{3}{2}, 0, -3)$	$\mathbf{p}_{02} = (\frac{3}{2}, 0, 3)$	$\mathbf{p}_{03} = (3, 0, 0)$
	$\mathbf{p}_{10} = (0, \frac{3}{2}, -3)$	$\mathbf{p}_{11} = (\frac{4}{3}, \frac{4}{3}, 0)$	$\mathbf{p}_{12} = (\frac{7}{6}, \frac{5}{3}, -\frac{10}{3})$	$\mathbf{p}_{13} = (3, 1, 1)$
	$\mathbf{p}_{20} = (0, \frac{3}{2}, 3)$	$\mathbf{p}_{21} = (\frac{5}{3}, \frac{7}{6}, \frac{10}{3})$	$\mathbf{p}_{22} = (\frac{11}{6}, \frac{11}{6}, 0)$	$\mathbf{p}_{23} = (3, 2, 1)$
S_2	$\mathbf{p}_{00} = (0, \frac{9}{8}, -\frac{3}{4})$	$\mathbf{p}_{01} = (\frac{7}{16}, \frac{49}{40}, -\frac{1}{5})$	$\mathbf{p}_{02} = (\frac{31}{40}, \frac{13}{10}, \frac{11}{16})$	$\mathbf{p}_{03} = (\frac{137}{128}, \frac{879}{640}, \frac{187}{160})$
	$\mathbf{p}_{04} = (\frac{11}{8}, \frac{29}{20}, \frac{3}{2})$	$\mathbf{p}_{05} = (\frac{27}{16}, \frac{25}{16}, \frac{7}{4})$	$\mathbf{p}_{10} = (\frac{7}{16}, \frac{37}{40}, -\frac{4}{5})$	$\mathbf{p}_{11} = (\frac{157}{200}, \frac{217}{200}, -\frac{27}{50})$
	$\mathbf{p}_{12} = (\frac{3367}{3200}, \frac{3901}{3200}, \frac{49}{800})$	$\mathbf{p}_{13} = (\frac{2073}{1600}, \frac{431}{320}, \frac{13}{40})$	$\mathbf{p}_{14} = (\frac{1251}{800}, \frac{1177}{800}, \frac{21}{40})$	$\mathbf{p}_{15} = (\frac{37}{20}, \frac{13}{8}, \frac{7}{10})$
	$\mathbf{p}_{20} = (\frac{3}{4}, \frac{3}{4}, -\frac{81}{80})$	$\mathbf{p}_{21} = (\frac{3287}{3200}, \frac{3053}{3200}, -\frac{679}{800})$	$\mathbf{p}_{22} = (\frac{199}{160}, \frac{181}{160}, -\frac{163}{400})$	$\mathbf{p}_{23} = (\frac{4659}{3200}, \frac{4149}{3200}, -\frac{223}{800})$
	$\mathbf{p}_{24} = (\frac{171}{100}, \frac{29}{20}, -\frac{59}{400})$	$\mathbf{p}_{25} = (\frac{1277}{640}, \frac{1039}{640}, \frac{3}{160})$	$\mathbf{p}_{30} = (\frac{637}{640}, \frac{351}{640}, -\frac{189}{160})$	$\mathbf{p}_{31} = (\frac{1957}{1600}, \frac{251}{320}, -\frac{197}{200})$
	$\mathbf{p}_{32} = (\frac{903}{640}, \frac{3173}{3200}, -\frac{99}{160})$	$\mathbf{p}_{33} = (\frac{517}{320}, \frac{1887}{1600}, -\frac{14}{25})$	$\mathbf{p}_{34} = (\frac{6023}{3200}, \frac{4301}{3200}, -\frac{343}{800})$	$\mathbf{p}_{35} = (\frac{2703}{320}, \frac{97}{64}, -\frac{7}{40})$
	$\mathbf{p}_{40} = (\frac{99}{80}, \frac{23}{80}, -\frac{3}{4})$	$\mathbf{p}_{41} = (\frac{1149}{800}, \frac{439}{800}, -\frac{101}{200})$	$\mathbf{p}_{42} = (\frac{647}{400}, \frac{311}{400}, -\frac{87}{400})$	$\mathbf{p}_{43} = (\frac{5887}{3200}, \frac{125}{128}, -\frac{183}{800})$
	$\mathbf{p}_{44} = (\frac{859}{400}, \frac{91}{80}, -\frac{7}{100})$	$\mathbf{p}_{45} = (\frac{101}{40}, \frac{103}{80}, \frac{7}{20})$	$\mathbf{p}_{50} = (\frac{3}{2}, 0, 0)$	$\mathbf{p}_{51} = (\frac{27}{16}, \frac{23}{80}, \frac{3}{20})$
	$\mathbf{p}_{52} = (\frac{241}{128}, \frac{343}{640}, \frac{7}{32})$	$\mathbf{p}_{53} = (\frac{687}{320}, \frac{237}{320}, \frac{1}{40})$	$\mathbf{p}_{54} = (\frac{101}{40}, \frac{71}{80}, \frac{3}{20})$	$\mathbf{p}_{55} = (3, 1, \frac{3}{4})$

Example 3. We consider the irreducible surfaces S_1 and S_2 of degree (2, 3) and (5, 5), respectively. Their control points are shown in Table 2. This example matches Case 1 of Lemma 4, and hence system (22) has a unique solution $(a_1, a_2, b_1, b_2, c_1, c_2, d_1, d_2) = (0, 1/2, 1/2, 0, 1, 1/2, 1/2, 1)$. Surfaces S_1 and S_3 are shown in Fig. 4a. and Fig. 4b., respectively. They have coincidence and their coincident part is surface S_2 , see Fig. 4c.

Example 4. We test for coincidence the irreducible surfaces S_1 and S_2 of degree (2, 2) and (4, 4), respectively. Their control points are shown in Table 3. The corresponding vectors $\rho, \rho^{1,0}, \rho^{0,1}$ are coplanar and hence, matrix M , defined by (25) has rank 2. We apply Algorithm 2 and obtain the following two solutions $(a_1, a_2, b_1, b_2, c_1, c_2, d_1, d_2)$ to system (23),

$$\begin{aligned} &(-1, 1/3, 4/3, 1/2, 3/4, 3/2, -1/4, 13/12), \\ &(34/71, 271/183, 75/284, -181/549, -291/284, 15/122, -277/568, 329/366). \end{aligned}$$

The first of these solutions generates TPB surface whose control polygon coincides with the control polygon of S_2 . Hence, S_1 and S_2 have coincidence. Surfaces S_1, S_2 , and their coincident part S are shown in Fig. 5 The control points of S are shown in Table 3.




Figure 5: The irreducible TPB surfaces S_1 of degree (2, 2) and S_2 of degree (4, 4) and their coincident part S from Example 4. The corresponding control points are shown in Table 3. **a.** Surface S_1 ; **b.** Surface S_2 ; **c.** Surfaces S_1, S_2 , and S which is shown with its control polygon.

Example 5. In our final example we test for coincidence the irreducible surfaces S_1 and S_2 of degree (2, 4) and (6, 6), respectively. Their control points are shown in Table 4. Surfaces S_1 and S_2 are shown in Fig. 6a. and Fig. 6b.. The corresponding system (22) has a unique solution $(a_1, a_2, b_1, b_2, c_1, c_2, d_1, d_2) = (1/2, -1/2, 0, 1/3, 1/2, 7/6, -1, 2/3)$. The shared domain of S_1 and S_2 is a hexagon. We represent the coincident part S as a union of two TPB surfaces and compute their control points using blossoming, see Fig 6c.

4. Conclusions and future work

In this paper we considered and solved the problem for testing TPB surfaces for coincidence. We presented two different methods and develop two algorithms based on these methods that test two irreducible TPB surfaces for

Table 3: The control points of irreducible TPB surfaces of degree (2, 2) and (4, 4) from Example 4

S_1	$\mathbf{p}_{00} = (0, 0, 0)$	$\mathbf{p}_{01} = (1, 0, 1)$	$\mathbf{p}_{02} = (2, 0, \frac{3}{4})$	
	$\mathbf{p}_{10} = (0, \frac{7}{10}, -\frac{1}{2})$	$\mathbf{p}_{11} = (1, 1, 0)$	$\mathbf{p}_{12} = (2, 1, \frac{6}{5})$	
	$\mathbf{p}_{20} = (0, 2, -\frac{4}{3})$	$\mathbf{p}_{21} = (1, 1, 1)$	$\mathbf{p}_{22} = (2, 2, \frac{3}{4})$	
S_2	$\mathbf{p}_{00} = (\frac{2}{3}, -\frac{86}{45}, \frac{95}{36})$	$\mathbf{p}_{01} = (\frac{25}{24}, -\frac{35}{18}, \frac{409}{180})$	$\mathbf{p}_{02} = (\frac{17}{12}, -\frac{121}{90}, \frac{5951}{5760})$	$\mathbf{p}_{03} = (\frac{43}{24}, -\frac{10109}{11520}, \frac{8609}{23040})$
	$\mathbf{p}_{04} = (\frac{13}{6}, -\frac{1123}{2304}, \frac{5957}{23040})$	$\mathbf{p}_{10} = (\frac{3}{4}, -\frac{85}{108}, \frac{2027}{2160})$	$\mathbf{p}_{11} = (\frac{37}{32}, -\frac{93}{160}, \frac{9631}{11520})$	$\mathbf{p}_{12} = (\frac{25}{16}, -\frac{5743}{17280}, \frac{9151}{17280})$
	$\mathbf{p}_{13} = (\frac{63}{32}, -\frac{23423}{138240}, \frac{116701}{276480})$	$\mathbf{p}_{14} = (\frac{19}{8}, \frac{53}{7680}, \frac{23203}{46080})$	$\mathbf{p}_{20} = (\frac{5}{6}, \frac{1051}{2430}, -\frac{1273}{38880})$	$\mathbf{p}_{21} = (\frac{61}{48}, \frac{87029}{155520}, \frac{89543}{311040})$
	$\mathbf{p}_{22} = (\frac{41}{24}, \frac{427381}{933120}, \frac{1410439}{1866240})$	$\mathbf{p}_{23} = (\frac{103}{48}, \frac{3137}{7680}, \frac{5157}{5120})$	$\mathbf{p}_{24} = (\frac{31}{12}, \frac{1151}{2560}, \frac{154297}{138240})$	$\mathbf{p}_{30} = (\frac{11}{12}, \frac{317}{240}, \frac{101}{2160})$
	$\mathbf{p}_{31} = (\frac{133}{96}, \frac{5191}{4320}, \frac{22847}{34560})$	$\mathbf{p}_{32} = (\frac{89}{48}, \frac{108571}{103680}, \frac{18113}{13824})$	$\mathbf{p}_{33} = (\frac{223}{96}, \frac{5179}{5120}, \frac{150163}{92160})$	$\mathbf{p}_{34} = (\frac{67}{24}, \frac{259}{256}, \frac{12917}{7680})$
	$\mathbf{p}_{40} = (1, \frac{83}{45}, \frac{63}{80})$	$\mathbf{p}_{41} = (\frac{3}{2}, \frac{947}{576}, \frac{5461}{5760})$	$\mathbf{p}_{42} = (2, \frac{12431}{5760}, \frac{18029}{34560})$	$\mathbf{p}_{43} = (\frac{5}{2}, \frac{4631}{1920}, \frac{2587}{3840})$
	$\mathbf{p}_{44} = (3, \frac{741}{320}, \frac{621}{640})$			
S	$\mathbf{p}_{00} = (\frac{17}{21}, 0, \frac{4267}{7056})$	$\mathbf{p}_{01} = (\frac{31}{28}, 0, \frac{10609}{14112})$	$\mathbf{p}_{02} = (\frac{59}{42}, 0, \frac{34927}{42336})$	$\mathbf{p}_{03} = (\frac{143}{84}, 0, \frac{277}{336})$
	$\mathbf{p}_{04} = (2, 0, \frac{3}{4})$	$\mathbf{p}_{10} = (\frac{71}{84}, \frac{1051}{2352}, \frac{11633}{35280})$	$\mathbf{p}_{11} = (\frac{127}{112}, \frac{2227}{4704}, \frac{71797}{141120})$	$\mathbf{p}_{12} = (\frac{239}{168}, \frac{6931}{14112}, \frac{143093}{211680})$
	$\mathbf{p}_{13} = (\frac{575}{336}, \frac{1}{2}, \frac{5587}{6720})$	$\mathbf{p}_{14} = (2, \frac{1}{2}, \frac{39}{40})$	$\mathbf{p}_{20} = (\frac{37}{42}, \frac{4537}{5292}, \frac{9115}{42336})$	$\mathbf{p}_{21} = (\frac{65}{56}, \frac{1163}{1323}, \frac{182957}{423360})$
	$\mathbf{p}_{22} = (\frac{121}{84}, \frac{28897}{31752}, \frac{817367}{1270080})$	$\mathbf{p}_{23} = (\frac{289}{168}, \frac{479}{504}, \frac{8563}{10080})$	$\mathbf{p}_{24} = (2, 1, \frac{21}{20})$	$\mathbf{p}_{30} = (\frac{11}{12}, \frac{109}{90}, \frac{103}{392})$
	$\mathbf{p}_{31} = (\frac{19}{16}, \frac{6127}{5040}, \frac{4673}{9408})$	$\mathbf{p}_{32} = (\frac{35}{24}, \frac{4783}{3780}, \frac{24463}{35280})$	$\mathbf{p}_{33} = (\frac{83}{48}, \frac{457}{336}, \frac{5731}{6720})$	$\mathbf{p}_{34} = (2, \frac{3}{2}, \frac{39}{40})$
	$\mathbf{p}_{40} = (\frac{20}{21}, \frac{662}{441}, \frac{991}{2205})$	$\mathbf{p}_{41} = (\frac{17}{14}, \frac{1313}{882}, \frac{5867}{8820})$	$\mathbf{p}_{42} = (\frac{31}{21}, \frac{2074}{1323}, \frac{41659}{52920})$	$\mathbf{p}_{43} = (\frac{73}{42}, \frac{73}{42}, \frac{137}{168})$
	$\mathbf{p}_{44} = (2, 2, \frac{3}{4})$			




Figure 6: **a.** Irreducible TPB surfaces S_1 of degree (2, 4) and S_2 of degree (6, 6) from Example 5. The control points of S_1 and S_2 are shown in Table 3. **a.** S_1 ; **b.** S_1 and S_2 with its control polygon; **c.** The coincident part S is presented as two TPB surfaces.

Table 4: The control points of irreducible TPB surfaces of degree (2, 4) and (6, 6) from Example 5

S_1	$\mathbf{p}_{00} = (0, 0, 0)$	$\mathbf{p}_{01} = (0, \frac{7}{10}, -\frac{1}{2})$	$\mathbf{p}_{02} = (0, 2, -\frac{4}{5})$	$\mathbf{p}_{03} = (0, \frac{23}{10}, -\frac{4}{5})$
	$\mathbf{p}_{04} = (0, 3, 1)$	$\mathbf{p}_{10} = (1, 0, 1)$	$\mathbf{p}_{11} = (1, 1, 0)$	$\mathbf{p}_{12} = (1, 1, 1)$
	$\mathbf{p}_{13} = (1, \frac{6}{5}, \frac{6}{5})$	$\mathbf{p}_{14} = (1, \frac{3}{2}, \frac{4}{3})$	$\mathbf{p}_{20} = (2, 0, \frac{3}{4})$	$\mathbf{p}_{21} = (2, 1, \frac{6}{5})$
	$\mathbf{p}_{22} = (2, 2, \frac{3}{4})$	$\mathbf{p}_{23} = (2, 3, 1)$	$\mathbf{p}_{24} = (2, 4, \frac{3}{2})$	
S_2	$\mathbf{p}_{00} = (1, -\frac{79}{32}, \frac{2717}{768})$	$\mathbf{p}_{01} = (\frac{7}{6}, -\frac{4151}{2880}, \frac{19667}{17280})$	$\mathbf{p}_{02} = (\frac{4}{3}, -\frac{277}{1800}, \frac{4481}{14400})$	$\mathbf{p}_{03} = (\frac{3}{2}, \frac{1361}{3600}, \frac{44467}{64800})$
	$\mathbf{p}_{04} = (\frac{5}{3}, \frac{5447}{6075}, \frac{255913}{291600})$	$\mathbf{p}_{05} = (\frac{11}{6}, \frac{1991}{1215}, \frac{26399}{29160})$	$\mathbf{p}_{06} = (2, \frac{8}{3}, \frac{1687}{1620})$	$\mathbf{p}_{10} = (\frac{5}{6}, -\frac{3973}{2880}, \frac{4081}{1728})$
	$\mathbf{p}_{11} = (1, -\frac{3143}{4320}, \frac{191717}{207360})$	$\mathbf{p}_{12} = (\frac{7}{6}, \frac{2489}{8640}, \frac{250213}{777600})$	$\mathbf{p}_{13} = (\frac{4}{3}, \frac{9049}{12150}, \frac{1379747}{2332800})$	$\mathbf{p}_{14} = (\frac{3}{2}, \frac{175433}{145800}, \frac{668537}{874800})$
	$\mathbf{p}_{15} = (\frac{5}{3}, \frac{13499}{7290}, \frac{307297}{349920})$	$\mathbf{p}_{16} = (\frac{11}{6}, \frac{3341}{1215}, \frac{32099}{29160})$	$\mathbf{p}_{20} = (\frac{2}{3}, -\frac{2269}{3600}, \frac{22961}{21600})$	$\mathbf{p}_{21} = (\frac{5}{6}, -\frac{1883}{14400}, \frac{157141}{388800})$
	$\mathbf{p}_{22} = (1, \frac{26249}{38880}, \frac{2976581}{23328000})$	$\mathbf{p}_{23} = (\frac{7}{6}, \frac{2027447}{1944000}, \frac{1061101}{2332800})$	$\mathbf{p}_{24} = (\frac{4}{3}, \frac{520417}{364500}, \frac{6107587}{8748000})$	$\mathbf{p}_{25} = (\frac{3}{2}, \frac{96541}{48600}, \frac{264529}{291600})$
	$\mathbf{p}_{26} = (\frac{5}{3}, \frac{5609}{2025}, \frac{115331}{97200})$	$\mathbf{p}_{30} = (\frac{1}{2}, -\frac{203}{900}, \frac{14257}{32400})$	$\mathbf{p}_{31} = (\frac{2}{3}, \frac{55237}{194400}, \frac{96383}{583200})$	$\mathbf{p}_{32} = (\frac{5}{6}, \frac{1863737}{1944000}, \frac{17473}{233280})$
	$\mathbf{p}_{33} = (1, \frac{272407}{216000}, \frac{4416611}{10368000})$	$\mathbf{p}_{34} = (\frac{7}{6}, \frac{3084847}{1944000}, \frac{1652957}{2332800})$	$\mathbf{p}_{35} = (\frac{4}{3}, \frac{33577}{16200}, \frac{27907}{28800})$	$\mathbf{p}_{36} = (\frac{3}{2}, \frac{89399}{32400}, \frac{246491}{194400})$
	$\mathbf{p}_{40} = (\frac{1}{3}, \frac{1819}{12150}, \frac{18443}{145800})$	$\mathbf{p}_{41} = (\frac{1}{2}, \frac{12494}{18225}, -\frac{1339}{437400})$	$\mathbf{p}_{42} = (\frac{2}{3}, \frac{904483}{729000}, \frac{35749}{4374000})$	$\mathbf{p}_{43} = (\frac{5}{6}, \frac{318233}{216000}, \frac{29029}{77760})$
	$\mathbf{p}_{44} = (1, \frac{1014119}{583200}, \frac{48464383}{69984000})$	$\mathbf{p}_{45} = (\frac{7}{6}, \frac{100631}{46656}, \frac{6996317}{6998400})$	$\mathbf{p}_{46} = (\frac{4}{3}, \frac{133901}{48600}, \frac{1543601}{1166400})$	$\mathbf{p}_{50} = (\frac{1}{6}, \frac{2891}{4860}, -\frac{1411}{7290})$
	$\mathbf{p}_{51} = (\frac{1}{3}, \frac{1613}{1458}, -\frac{42221}{174960})$	$\mathbf{p}_{52} = (\frac{1}{2}, \frac{37217}{24300}, -\frac{18343}{145800})$	$\mathbf{p}_{53} = (\frac{2}{3}, \frac{328277}{194400}, \frac{169903}{583200})$	$\mathbf{p}_{54} = (\frac{5}{6}, \frac{2213237}{1166400}, \frac{2383159}{3499200})$
	$\mathbf{p}_{55} = (1, \frac{87773}{38880}, \frac{1983853}{1866240})$	$\mathbf{p}_{56} = (\frac{7}{6}, \frac{8009}{2880}, \frac{224443}{155520})$	$\mathbf{p}_{60} = (0, \frac{17}{15}, -\frac{203}{405})$	$\mathbf{p}_{61} = (0, \frac{17}{15}, -\frac{203}{405})$
	$\mathbf{p}_{62} = (\frac{1}{3}, \frac{22679}{12150}, -\frac{24437}{145800})$	$\mathbf{p}_{63} = (\frac{1}{2}, \frac{31721}{16200}, \frac{34481}{97200})$	$\mathbf{p}_{64} = (\frac{2}{3}, \frac{206657}{97200}, \frac{499307}{583200})$	$\mathbf{p}_{65} = (\frac{5}{6}, \frac{190249}{77760}, \frac{62737}{46656})$
	$\mathbf{p}_{66} = (1, \frac{7553}{2592}, \frac{112733}{62208})$			

coincidence and report their coincident part if it is present. The first algorithm works for surfaces of same degree (m, n) and the second one - for surfaces of degree (m, n) and $(m + n, m + n)$, respectively. We presented numerical experiments and gave examples to visualize and support the obtained results. Our next task for future research is to develop and implement an algorithm for testing TB surfaces for coincidence.

Acknowledgments

This work was partially supported by Sofia University Science Fund Grant No. 80-10-103/2023.

References

- Barton, D., Zippel, R., 1985. Polynomial decomposition algorithms. *J. Symbolic Comput.* 1, 159–168. doi:[10.1016/S0747-7171\(85\)80012-2](https://doi.org/10.1016/S0747-7171(85)80012-2).
- Berry, T., Patterson, R., 1997. The uniqueness of Bézier control points. *Comput. Aided Geom. Des.* 14, 877–879. doi:[10.1016/S0167-8396\(97\)00016-2](https://doi.org/10.1016/S0167-8396(97)00016-2).
- Chen, X.D., Ma, W., 2015. Rebuttal to “comment on the ‘coincidence condition of two Bézier curves of an arbitrary degree’ ”. *Comput. Graph.* 53, 167–169. doi:<https://doi.org/10.1016/j.cag.2015.10.007>.
- Chen, X.D., W. Ma, W., Deng, C., 2013. Conditions for the coincidence of two quartic Bézier curves. *Appl. Math. and Comput.* 225, 731–736. doi:[10.1016/j.amc.2013.09.053](https://doi.org/10.1016/j.amc.2013.09.053).
- Chen, X.D., Yang, C., Ma, W., 2016. Coincidence condition of two Bézier curves of an arbitrary degree. *Comput. Graph.* 54, 121–126. doi:[10.1016/j.cag.2015.07.013](https://doi.org/10.1016/j.cag.2015.07.013).
- Denker, W., Herron, G., 1997. Generalizing rational degree elevation. *Comput. Aided Geom. Des.* 14, 399–406. doi:[10.1016/S0167-8396\(96\)00036-2](https://doi.org/10.1016/S0167-8396(96)00036-2).
- Farin, G., 2002. *Curves and Surfaces for CAGD: A Practical Guide*. 5th ed., Morgan-Kaufmann, San Francisco.
- Farouki, R., 2012. The Bernstein polynomial basis: a centennial retrospective. *Comput. Aided Geom. Des.* 29, 379–419. doi:[10.1016/j.cagd.2012.03.001](https://doi.org/10.1016/j.cagd.2012.03.001).
- von zur Gathen, J., 1990. Functional decomposition of polynomials: the tame case. *J. Symbolic Comput.* 9, 281–299. doi:[10.1016/S0747-7171\(08\)80014-4](https://doi.org/10.1016/S0747-7171(08)80014-4).
- Goldman, R., 2003. *Pyramid algorithms: A dynamic programming approach to curves and surfaces for geometric modeling*. Elsevier Science (USA).
- Kozen, D., Landau, S., 1989. Polynomial decomposition algorithms. *J. Symbolic Comput.* 7, 445–456. doi:[https://doi.org/10.1016/S0747-7171\(89\)80027-6](https://doi.org/10.1016/S0747-7171(89)80027-6).
- Pekerman, D., Seong, J.K., Elber, G., Kim, M.S., 2005. Are two curves the same? *Comput.-Aided Des. Appl.* 2, 85–94. doi:[10.1080/16864360.2005.10738356](https://doi.org/10.1080/16864360.2005.10738356).
- Sánchez-Reyes, J., 2011. On the conditions for the coincidence of two cubic Bézier curves. *J. of Comput. Appl. Math.* 236, 1675–1677. doi:[10.1016/j.cam.2011.08.024](https://doi.org/10.1016/j.cam.2011.08.024).
- Sánchez-Reyes, J., 2014. The conditions for the coincidence or overlapping of two Bézier curves. *Appl. Math. Comput.* 248, 625–630. doi:[10.1016/j.amc.2014.10.008](https://doi.org/10.1016/j.amc.2014.10.008).
- Sánchez-Reyes, J., 2015a. Comment on the “coincidence condition of two Bézier curves of an arbitrary degree”. *Comput. Graph.* 53, 166. doi:[10.1016/j.cag.2015.10.006](https://doi.org/10.1016/j.cag.2015.10.006).
- Sánchez-Reyes, J., 2015b. Detecting symmetries in polynomial Bézier curves. *J. of Comput. Appl. Math.* 288, 274–283. doi:[10.1016/j.cam.2015.04.025](https://doi.org/10.1016/j.cam.2015.04.025).
- Vlachkova, K., 2017. Comparing Bézier curves and surfaces for coincidence, in: Georgiev, K., Todorov, M., Georgiev, I. (Eds.), *Advanced Computing in Industrial Mathematics*. Springer International Publishing. volume 681 of *LNCS, Studies in Computational Intelligence*, pp. 239–250. doi:[10.1007/978-3-319-49544-6_20](https://doi.org/10.1007/978-3-319-49544-6_20).
- W.-K. Wang, W.K., Zhang, H., Liu, X.M., Paul, J.C., 2011. Conditions for coincidence of two cubic Bézier curves. *J. Comput. Appl. Math.* 235, 5198–5202. doi:[10.1016/j.cam.2011.05.006](https://doi.org/10.1016/j.cam.2011.05.006).
- Yang, L.Q., Zeng, X.M., 2008. Trimming Bézier surfaces on Bézier surfaces via blossoming, in: Chen, F., Jüttler, B. (Eds.), *LNCS 4975*, Springer, Berlin Heidelberg. pp. 578–584. doi:[10.1007/978-3-540-79246-8_50](https://doi.org/10.1007/978-3-540-79246-8_50).