

NATURAL WEIGHTS FOR UNIFORM APPROXIMATION BY THE SZÁSZ-MIRAKJAN OPERATOR

BORISLAV R. DRAGANOV AND KAMEN G. IVANOV

ABSTRACT. We establish a sharp characterization of the error of the Szász-Mirakjan operator in uniform norm with power-type weights in terms of a K -functional. The weight exponents are optimal. We also state a sharp characterization of the K -functional by means of the classical unweighted fixed-step modulus of smoothness.

1. INTRODUCTION

1.1. **Notations and main result.** The Szász-Mirakjan operator for a function f defined on $[0, \infty)$ is given by

$$S_n f(x) = \sum_{k=0}^{\infty} f\left(\frac{k}{n}\right) s_{n,k}(x), \quad s_{n,k}(x) = e^{-nx} \frac{(nx)^k}{k!}, \quad n \geq 1, \quad x \geq 0.$$

Here n is not necessarily an integer.

In order to describe the approximation properties of S_n we need a number of function spaces. Let $C[0, \infty)$ denote the space of all continuous functions on $[0, \infty)$ and $L_{\infty}[0, \infty)$ denote the Lebesgue measurable and essentially bounded on $[0, \infty)$ functions with the essential supremum norm $\|\cdot\|$. For continuous functions $\|\cdot\|$ coincides with the uniform norm on $[0, \infty)$. Also, we denote the first derivative operator by $D = \frac{d}{dx}$, thus $Dg(x) = g'(x)$ and $D^2g(x) = g''(x)$. We further set $\varphi(x) = x$. Now, for a weight function w on $(0, \infty)$ we put

$$\begin{aligned} C(w)(0, \infty) &= \{f \in C[0, \infty) : wf \in L_{\infty}[0, \infty)\}, \\ W^2(w\varphi)(0, \infty) &= \{g \in AC_{loc}(0, \infty) : Dg \in AC_{loc}(0, \infty), w\varphi D^2g \in L_{\infty}[0, \infty)\}, \\ W^3(w\varphi^{3/2})(0, \infty) &= \{g \in AC_{loc}(0, \infty) : Dg, D^2g \in AC_{loc}(0, \infty), \\ &\quad w\varphi^{3/2} D^3g \in L_{\infty}[0, \infty)\}, \end{aligned}$$

where $AC_{loc}(0, \infty)$ consists of the functions which are absolutely continuous in $[a, b]$ for every $[a, b] \subset (0, \infty)$. Finally, π_1 stands for the set of algebraic polynomials of first degree.

In [13] the second author introduces the concept of a *natural weight* for an approximation operator.

2010 *Mathematics Subject Classification.* Primary 41A25; Secondary 41A15, 41A27.

Key words and phrases. Szász-Mirakjan operator, modulus of smoothness, K -functional, rate of convergence, weighted uniform approximation.

Both authors are supported by grant DDVU 02/30 of the Fund for Scientific Research of the Bulgarian Ministry of Education and Science.

Definition 1.1. A weight w is called a *natural weight* for approximation by a sequence Q_n of operators in a specified norm if the norm of the weighted approximation error $w(f - Q_n f)$ allows matching direct and strong inverse estimates for the widest reasonable class of functions f .

Here we shall consider weights on $(0, \infty)$ of the form

$$(1.1) \quad w(x) = w(\gamma_0, \gamma_\infty; x) = \left(\frac{x}{1+x} \right)^{\gamma_0} (1+x)^{\gamma_\infty}$$

and show that they are natural for the uniform approximation by the Szász-Mirakjan operator for any $\gamma_0 \in [-1, 0]$ and $\gamma_\infty \in \mathbb{R}$. More precisely, let the K -functional $K_w(f, t)$ be defined for $f \in C(w)(0, \infty) + \pi_1$ and $t > 0$ by

$$(1.2) \quad K_w(f, t) = \inf \{ \|w(f - g)\| + t \|w\varphi D^2 g\| : g \in W^2(w\varphi)(0, \infty) \}.$$

We establish the following characterization.

Theorem 1.2. *Let $w = w(\gamma_0, \gamma_\infty)$ be given by (1.1) with $\gamma_0 \in [-1, 0]$ and $\gamma_\infty \in \mathbb{R}$. Then for all $f \in C(w)(0, \infty) + \pi_1$ and $n \geq 1$ there holds*

$$\|w(f - S_n f)\| \sim K_w \left(f, \frac{1}{n} \right).$$

The relation $\psi(f, t) \sim \theta(f, t)$ means that there exists a positive constant independent of f and t such that

$$c^{-1}\theta(f, t) \leq \psi(f, t) \leq c\theta(f, t).$$

Let us note that the range of γ_0 cannot be reasonably extended. Indeed, if $\gamma_0 < -1$ then we must assume that $f(x) = 0$ in a neighbourhood of 0, otherwise $S_n f$ will not be bounded. On the other hand, if $\gamma_0 > 0$, then $f(x)$ is not generally defined at 0 and hence $S_n f$ is not defined. Moreover, we cannot settle this case by restricting to functions $f \in C[0, \infty)$ because then S_n is not a bounded operator in the weighted uniform norm.

1.2. Characterization of the weighted K -functionals. The K -functional $K_w(f, t)$ is equivalent to the Ditzian-Totik modulus of smoothness if $\gamma_0 = 0$ (see [4, Theorem 6.1.1]), whereas for $\gamma_0 < 0$ it is only weakly equivalent to the so called weighted main-part modulus (see [4, Section 6.2]). Also, this K -functional with $\gamma_0 = \gamma_\infty = 0$ was shown to be equivalent to a modulus introduced by the second author in [12]. In a series of papers [5, 6, 7, 8] the authors introduced and studied moduli of smoothness which are equivalent to weighted K -functionals like $K_w(f, t)$. They are defined by means of the classical unweighted fixed-step modulus and a continuous linear transform of the function. In particular, we can characterize $K_w(f, t)$ by such moduli by first splitting the singularities

$$K_w(f, t) \sim K_w(f, t)_{[0,2]} + K_w(f, t)_{[1,\infty)},$$

which is valid for all $f \in C(w)(0, \infty)$ and $t \in (0, 1]$. The K -functional $K_w(f, t)_I$ is defined as in (1.2) with the supremum taken on the interval I . By [8, Theorem 6.2 and 6.8] we have

$$(1.3) \quad K_w(f, t)_{[0,2]} \sim \omega_2(\mathcal{A}_{\gamma_0} f, t)_{[0,2]}$$

where $\omega_2(f, t)_{[0,2]}$ is the classical modulus of smoothness of second order in uniform norm on the interval $[0, 2]$ and

$$\mathcal{A}_{\gamma_0} f(x) = \begin{cases} f(x^2) - x \int_1^x y^{-2} f(y^2) dy & \text{for } \gamma_0 = 0, \\ x^{2\gamma} f(x^2) + \frac{4\gamma^2 - 1}{5} x \int_1^x y^{2\gamma-2} f(y^2) dy \\ \quad - \frac{4(\gamma_0 + 2)(\gamma_0 + 3)}{5} x^{-4} \int_0^x y^{2\gamma+3} f(y^2) dy & \text{for } -1 < \gamma_0 < 0, \\ x^{-2} f(x^2) + 3x \int_1^x y^{-4} f(y^2) dy & \text{for } \gamma_0 = -1, \end{cases}$$

for $x \in [0, 2]$. Relation (1.3) holds provided that wf has a limit at 0, which is, moreover, equal to 0 if $\gamma_0 \in (-1, 0)$.

Similarly, one can construct a modulus equivalent to $K_w(f, t)_{[1, \infty)}$. We shall present this case in [9].

1.3. Related works. Many mathematicians investigated the approximation behaviour of the Szász-Mirakjan operator. First, Becker [2] established direct and weak converse estimates for $w(x) = (1+x)^{-N}$, $N \in \mathbb{N}$. Amonov [1] proved a weak equivalence result for $w(x) = (1+x)^\gamma$, $\gamma \leq 0$. He studied also more general weights. Holhoş [11] introduced a general approach to verifying direct estimates in weighted uniform norm. As an application he obtained a direct result for S_n in the case $w(x) = (1+x)^\gamma$, $\gamma \leq 0$.

As for strong converse estimates, Totik [15] established the assertion of Theorem 1.2 in the unweighted case $w = 1$. Quite recently Finta [10] developed a general method, which in particular enable him to verify a two-term strong converse inequality for the weights $w(x) = x^\gamma$, $\gamma \in (-1, 0]$.

As can be seen, most researchers were interested in studying the rate of approximation in weighted spaces that are larger than $C[0, \infty)$ and thus allowing for more function to be approximated. In our opinion, it is also worthwhile finding in which weighted spaces the norm of the approximation error can be neatly characterized since that reveals intrinsic properties of the operator.

2. PROOF OF THEOREM 1.2

First, let us recall several basic properties of the Szász-Mirakjan operator:

(2.1) S_n is a positive linear operator;

(2.2) $S_n f = f \quad \forall f \in \pi_1$;

(2.3) $S_n((\circ - x)^2)(x) = \frac{x}{n}$, $S_n((\circ - x)^3)(x) = \frac{x}{n^2}$;

(2.4) $S_n((\circ - x)^4)(x) = d_1 \frac{x^2}{n^2} + d_2 \frac{x}{n^3}$, $S_n((\circ - x)^6)(x) = d_3 \frac{x^3}{n^3} + d_4 \frac{x^2}{n^4} + d_5 \frac{x}{n^5}$;

(2.5) $S_n f = (S_1(f_{1/n}))_n$, where $F_\nu(x) = F(\nu x)$, $\nu > 0$.

Above, d_1, \dots, d_5 are constants independent of n . For (2.3)-(2.4) see e.g. [4, (9.1.13) and Lemma 9.5.5].

The proof of Theorem 1.2 is based on [3, Theorem 4.1 and (1.2)-(1.4)]. So, we proceed to the verification of their hypotheses. The first concerns the boundedness of the Szász-Mirakjan operator in weighted uniform norm. Below c, c_1, c_2, \dots denote positive constants which do not depend on f and n .

Proposition 2.1. *Let $w = w(\gamma_0, \gamma_\infty)$ be given by (1.1) with $\gamma_0 \in [-1, 0]$ and $\gamma_\infty \in \mathbb{R}$. Then*

$$\|wS_n f\| \leq c_1 \|wf\| \quad \forall f \in C(w)(0, \infty) \quad \forall n \geq 1.$$

Proof. Using the positivity of S_n , we get

$$|S_n f(x)| \leq (S_n(w^{-1}))(x) \|wf\|$$

Thus it remains to show that

$$(2.6) \quad (S_n(w^{-1}))(x) \leq cw(x)^{-1}.$$

To this end we apply a routine technique based on Hölder's inequality. If $\gamma_0 \neq \gamma_\infty$, we fix $r \in \mathbb{Z}$ such that $r/(\gamma_0 - \gamma_\infty) \geq 1$; if $\gamma_0 = \gamma_\infty$, we set $r = 1$. Then consecutive applications of Hölder's inequality yield

$$(2.7) \quad \begin{aligned} & \sum_{k \geq 0} \left(\frac{k}{n}\right)^{-\gamma_0} \left(1 + \frac{k}{n}\right)^{\gamma_0 - \gamma_\infty} s_{n,k}(x) \\ & \leq \left\{ \sum_{k \geq 0} \frac{k}{n} \left(1 + \frac{k}{n}\right)^{\gamma_0 - \gamma_\infty} s_{n,k}(x) \right\}^{-\gamma_0} \left\{ \sum_{k \geq 0} \left(1 + \frac{k}{n}\right)^{\gamma_0 - \gamma_\infty} s_{n,k}(x) \right\}^{1+\gamma_0} \\ & \leq \left\{ \sum_{k \geq 0} \frac{k}{n} \left(1 + \frac{k}{n}\right)^r s_{n,k}(x) \right\}^{\gamma_0(\gamma_\infty - \gamma_0)/r} \left\{ \sum_{k \geq 0} \frac{k}{n} s_{n,k}(x) \right\}^{-\gamma_0(1+(\gamma_\infty - \gamma_0)/r)} \\ & \quad \times \left\{ \sum_{k \geq 0} \left(1 + \frac{k}{n}\right)^r s_{n,k}(x) \right\}^{(1+\gamma_0)(\gamma_0 - \gamma_\infty)/r} \left\{ \sum_{k \geq 0} s_{n,k}(x) \right\}^{(1+\gamma_0)(1+(\gamma_\infty - \gamma_0)/r)}. \end{aligned}$$

It is known that (cf. [4, p. 163]),

$$(2.8) \quad \sum_{k \geq 0} \left(1 + \frac{k}{n}\right)^r s_{n,k}(x) \leq c(1+x)^r, \quad r \in \mathbb{Z};$$

hence also

$$(2.9) \quad \sum_{k \geq 0} \frac{k}{n} \left(1 + \frac{k}{n}\right)^r s_{n,k}(x) \leq cx \sum_{k \geq 0} \left(1 + \frac{k}{n}\right)^r s_{n,k}(x) \leq cx(1+x)^r.$$

Now, (2.6) readily follows from (2.2) and (2.7)-(2.9). \square

Next, we establish a Jackson-type inequality.

Proposition 2.2. *Let $w = w(\gamma_0, \gamma_\infty)$ be given by (1.1) with $\gamma_0 \in [-1, 0]$ and $\gamma_\infty \in \mathbb{R}$. Then*

$$\|w(g - S_n g)\| \leq \frac{c_2}{n} \|w\varphi D^2 g\| \quad \forall g \in W^2(w\varphi)(0, \infty) \quad \forall n \geq 1.$$

Proof. Since $g \in W^2(w\varphi)(0, \infty)$, then $g \in C[0, \infty)$. By Taylor's formula we have for $x > 0$ and $t \geq 0$ that

$$g(t) = g(x) + (t-x)Dg(x) + \int_x^t (t-u)D^2g(u) du.$$

We apply S_n to both sides of this identity with regard to the variable t . Then, taking into account that S_n preserves the linear functions (see (2.2)), we get

$$S_n g(x) = g(x) + S_n \left(\int_x^\circ (\circ - u) D^2 g(u) du \right) (x).$$

Next, by the positivity of S_n , we derive the estimate

$$(2.10) \quad \begin{aligned} |g(x) - S_n g(x)| &\leq S_n \left(\int_x^\circ (\circ - u) |D^2 g(u)| du \right) (x) \\ &\leq R_{1,n}(x) \|w\varphi D^2 g\|, \end{aligned}$$

where we have set

$$R_{1,n}(w; x) = S_n \left(\int_0^\infty Q_1(\circ, u, x) w(u)^{-1} du \right) (x)$$

and $Q_1(t, u, x) = |t - u|u^{-1}$ for u between t and x and $Q_1(t, u, x) = 0$ otherwise.

Thus it is sufficient to show that

$$(2.11) \quad R_{1,n}(w; x) \leq \frac{c}{n} w(x)^{-1}.$$

Just as in the proof of Proposition 2.1 we get

$$(2.12) \quad \begin{aligned} R_{1,n}(w(\gamma_0, \gamma_\infty); x) &\leq R_{1,n}(w(-1, -r-1); x)^{\gamma_0(\gamma_\infty - \gamma_0)/r} \\ &\quad \times R_{1,n}(w(-1, 0); x)^{-\gamma_0(1 + (\gamma_\infty - \gamma_0)/r)} \\ &\quad \times R_{1,n}(w(0, -r); x)^{(1 + \gamma_0)(\gamma_0 - \gamma_\infty)/r} \\ &\quad \times R_{1,n}(w(0, 0); x)^{(1 + \gamma_0)(1 + (\gamma_\infty - \gamma_0)/r)}. \end{aligned}$$

We estimate separately each of the $R_{1,n}$'s above. By (2.3) we immediately get

$$(2.13) \quad R_{1,n}(w(-1, 0); x) \leq S_n((\circ - x)^2)(x) = \frac{x}{n}.$$

Next, we observe that for u between t and x there hold

$$(2.14) \quad \frac{|t - u|}{u} \leq \frac{|t - x|}{x}, \quad (1 + u)^r \leq (1 + x)^r + (1 + t)^r.$$

The first of these inequalities is verified directly and the second one is trivial. The first inequality above and (2.2) yield

$$(2.15) \quad R_{1,n}(w(0, 0); x) \leq \frac{1}{x} S_n((\circ - x)^2)(x) = \frac{1}{n}.$$

Further, by means of the second inequality in (2.14) we arrive at

$$(2.16) \quad R_{1,n}(w(-1, -r-1); x) \leq (1 + x)^r S_n((\circ - x)^2)(x) + S_n((\circ - x)^2(1 + \circ)^r)(x)$$

Let us consider $S_n((\circ - x)^2(1 + \circ)^r)(x)$. For $nx \geq 1$ we apply the Cauchy inequality, (2.4) and (2.8) to get

$$\begin{aligned} S_n((\circ - x)^2(1 + \circ)^r)(x) &\leq \sqrt{S_n((\circ - x)^4)(x)} \sqrt{S_n((1 + \circ)^{2j})(x)} \\ &\leq \frac{cx(1 + x)^r}{n}. \end{aligned}$$

To verify this inequality for $nx \leq 1$ we proceed as follows:

$$S_n((\circ - x)^2(1 + \circ)^r)(x) = \sum_{k \geq 0} \left(\frac{k}{n} - x \right)^2 \left(1 + \frac{k}{n} \right)^r s_{n,k}(x)$$

$$\begin{aligned}
&= \sum_{k \geq 0} \left(\frac{k}{n}\right)^2 \left(1 + \frac{k}{n}\right)^r s_{n,k}(x) - 2x \sum_{k \geq 0} \frac{k}{n} \left(1 + \frac{k}{n}\right)^r s_{n,k}(x) \\
&\quad + x^2 \sum_{k \geq 0} \left(1 + \frac{k}{n}\right)^r s_{n,k}(x) \\
&\leq cx \sum_{k \geq 0} \frac{k}{n} \left(1 + \frac{k}{n}\right)^r s_{n,k}(x) + c \left(\frac{x}{n} + x^2\right) \sum_{k \geq 0} \left(1 + \frac{k}{n}\right)^r s_{n,k}(x) \\
&\leq c \left(\frac{x}{n} + x^2\right) (1+x)^r \leq \frac{cx(1+x)^r}{n},
\end{aligned}$$

as at the last but one step we have applied (2.8) and (2.9).

Thus we have established the estimate

$$(2.17) \quad S_n((\circ - x)^2(1 + \circ)^r)(x) \leq \frac{cx(1+x)^r}{n}, \quad x \geq 0.$$

Now, (2.16), (2.3) and (2.17) imply

$$(2.18) \quad R_{1,n}(w(-1, -r-1); x) \leq \frac{cx(1+x)^r}{n}.$$

Finally, by means of (2.14) we get

$$R_{1,n}(w(0, -r); x) \leq \frac{(1+x)^r}{x} S_n((\circ - x)^2)(x) + \frac{1}{x} S_n((\circ - x)^2(1 + \circ)^r)(x),$$

which along with (2.3) and (2.17), gives

$$(2.19) \quad R_{1,n}(w(0, -r); x) \leq \frac{c(1+x)^r}{n}, \quad x \geq 0.$$

The estimate (2.11) follows from (2.12), (2.13), (2.15), (2.18) and (2.19). The proof of the proposition is completed. \square

We shall also need the following Voronovskaya-type inequality.

Proposition 2.3. *Let $w = w(\gamma_0, \gamma_\infty)$ be given by (1.1) with $\gamma_0 \in [-1, 0]$ and $\gamma_\infty \in \mathbb{R}$. Then*

$$\left\| w \left(S_n g - g - \frac{1}{2n} \varphi D^2 g \right) \right\| \leq \frac{c_3}{n^{3/2}} \|w\varphi^{3/2} D^3 g\| \quad \forall g \in W^3(w\varphi^{3/2})(0, \infty) \quad \forall n \geq 1.$$

Proof. The proof is quite similar to the one of the previous proposition. We first use Taylor's formula to get the representation

$$g(t) = g(x) + (t-x)Dg(x) + \frac{(t-x)^2}{2} D^2 g(x) + \frac{1}{2} \int_x^t (t-u)^2 D^3 g(u) du.$$

We apply S_n to both sides of this identity with regard to the variable t . Then, taking into account (2.2) and (2.3), we get

$$S_n g(x) = g(x) + \frac{1}{2n} \varphi(x) D^2 g(x) + \frac{1}{2} S_n \left(\int_x^\circ (\circ - u)^2 D^3 g(u) du \right) (x).$$

Next, by the positivity of S_n , we derive the estimate

$$\begin{aligned}
(2.20) \quad \left| S_n g(x) - g(x) - \frac{1}{2n} \varphi(x) D^2 g(x) \right| &\leq \frac{1}{2} S_n \left(\left| \int_x^\circ (\circ - u)^2 |D^3 g(u)| du \right| \right) (x) \\
&\leq \frac{1}{2} R_{2,n}(x) \|w\varphi^{3/2} D^3 g\|,
\end{aligned}$$

where we have set

$$R_{2,n}(w; x) = S_n \left(\int_0^\infty Q_2(\circ, u, x) w(u)^{-1} du \right) (x)$$

and $Q_2(t, u, x) = (t-u)^2 u^{-3/2}$ for u between t and x and $Q_2(t, u, x) = 0$ otherwise.

We shall show that

$$(2.21) \quad R_{2,n}(w(\gamma_0, \gamma_\infty); x) \leq \frac{c}{n^{3/2}} w(x)^{-1}.$$

As in the proof of the previous propositions we get

$$\begin{aligned} R_{2,n}(w(\gamma_0, \gamma_\infty); x) &\leq R_{2,n}(w(-1, -r-1); x)^{\gamma_0(\gamma_\infty - \gamma_0)/r} \\ &\quad \times R_{2,n}(w(-1, 0); x)^{-\gamma_0(1+(\gamma_\infty - \gamma_0)/r)} \\ &\quad \times R_{2,n}(w(0, -r); x)^{(1+\gamma_0)(\gamma_0 - \gamma_\infty)/r} \\ &\quad \times R_{2,n}(w(0, 0); x)^{(1+\gamma_0)(1+(\gamma_\infty - \gamma_0)/r)}. \end{aligned}$$

Thus it is sufficient to establish (2.21) only for each of the $R_{2,n}$'s on the right above. This is done by arguments similar to those in the proof of the previous proposition based on the following relations:

$$(2.22) \quad \frac{(t-u)^2}{u^{1/2}} \leq \frac{(t-x)^2}{x^{1/2}}, \quad \frac{(t-u)^2}{u^{3/2}} \leq \frac{(t-x)^2}{x^{3/2}} \quad \text{for } u \text{ between } t \text{ and } x,$$

$$(2.23) \quad S_n(|\circ - x|^3(1+\circ)^i)(x) \leq \frac{c}{n^{3/2}} x^{3/2}(1+x)^i, \quad i \in \mathbb{Z}, \quad nx \geq 1,$$

$$(2.24) \quad \sum_{k \geq 2} \left(\frac{k}{n} - x \right)^3 \left(1 + \frac{k}{n} \right)^i s_{n,k}(x) \leq \frac{c}{n^{3/2}} x^{3/2}(1+x)^i, \quad i \in \mathbb{Z}, \quad nx \leq 1,$$

and

$$(2.25) \quad \left| \int_x^{k/n} \frac{\left(\frac{k}{n} - u \right)^2}{u^{3/2}} u^\alpha (1+u)^\beta du \right| s_{n,k}(x) \leq \frac{c}{n^{3/2}} w(-\alpha, -\beta; x)^{-1}, \quad nx \leq 1,$$

for $k = 0, 1$ and each of the couples

$$(\alpha, \beta) = (0, 0), \quad (\alpha, \beta) = (1, 0), \quad (\alpha, \beta) = (0, r), \quad (\alpha, \beta) = (1, r).$$

The inequalities (2.22) are verified directly.

To prove (2.23) we apply the Cauchy inequality, (2.4) and (2.8) and thus get

$$\begin{aligned} S_n(|\circ - x|^3(1+\circ)^i)(x) &\leq \sqrt{S_n((\circ - x)^6)(x)} \sqrt{S_n((1+\circ)^{2i})(x)} \\ &\leq \frac{c}{n^{3/2}} x^{3/2}(1+x)^i \end{aligned}$$

since $nx \geq 1$.

To establish (2.24) we observe that

$$\begin{aligned}
& \sum_{k \geq 2} \left(\frac{k}{n} - x \right)^3 \left(1 + \frac{k}{n} \right)^i s_{n,k}(x) \\
&= \sum_{k \geq 2} \frac{k}{n} \left(\frac{k}{n} - x \right)^2 \left(1 + \frac{k}{n} \right)^i s_{n,k}(x) - x \sum_{k \geq 2} \left(\frac{k}{n} - x \right)^2 \left(1 + \frac{k}{n} \right)^i s_{n,k}(x) \\
&\leq cx \sum_{k \geq 1} \left(\frac{k+1}{n} - x \right)^2 \left(1 + \frac{k}{n} \right)^i s_{n,k}(x) \\
&= cx \left[\sum_{k \geq 1} \left(\frac{k}{n} - x \right)^2 \left(1 + \frac{k}{n} \right)^i s_{n,k}(x) + \frac{2}{n} \sum_{k \geq 1} \left(\frac{k}{n} - x \right) \left(1 + \frac{k}{n} \right)^i s_{n,k}(x) \right. \\
&\quad \left. + \frac{1}{n^2} \sum_{k \geq 1} \left(1 + \frac{k}{n} \right)^i s_{n,k}(x) \right] \\
&\leq \frac{cx^2(1+x)^i}{n} \leq \frac{cx^{3/2}(1+x)^i}{n^{3/2}}, \quad nx \leq 1,
\end{aligned}$$

as at the last but one step we have used (2.8), (2.9), (2.17) and the estimate

$$\begin{aligned}
\frac{1}{n^2} \sum_{k \geq 1} \left(1 + \frac{k}{n} \right)^i s_{n,k}(x) &= \frac{x}{n} \sum_{k \geq 1} \frac{1}{k} \left(1 + \frac{k}{n} \right)^i s_{n,k-1}(x) \\
&\leq \frac{cx}{n} \sum_{k \geq 0} \left(1 + \frac{k}{n} \right)^i s_{n,k}(x) \leq \frac{cx(1+x)^i}{n}.
\end{aligned}$$

It remains to prove (2.25). Set

$$A_{n,k}(\alpha, \beta; x) = \left| \int_x^{k/n} \frac{\left(\frac{k}{n} - u \right)^2}{u^{3/2}} u^\alpha (1+u)^\beta du \right| s_{n,k}(x).$$

For $\alpha = \beta = 0$ we have by straightforward calculations

$$A_{n,0}(0, 0; x) = \int_0^x \frac{u^2}{u^{3/2}} du s_{n,0}(x) = \frac{2}{3} x^{3/2} e^{-nx} \leq \frac{c}{n^{3/2}} w(0, 0; x)^{-1}$$

and

$$\begin{aligned}
A_{n,1}(0, 0; x) &= \int_x^{1/n} \frac{\left(\frac{1}{n} - u \right)^2}{u^{3/2}} du s_{n,1}(x) \\
&= e^{-nx} \left(2(1-nx)^2 \frac{x^{1/2}}{n} - 4nx \int_x^{1/n} \frac{\frac{1}{n} - u}{u^{1/2}} du \right) \\
&\leq \frac{c}{n^{3/2}} w(0, 0; x)^{-1}.
\end{aligned}$$

For $\alpha = 1, \beta = 0$ we calculate

$$A_{n,0}(1, 0; x) = \int_0^x \frac{u^2}{u^{3/2}} u du s_{n,0}(x) = \frac{2}{5} x^{5/2} e^{-nx} \leq \frac{c}{n^{3/2}} w(-1, 0; x)^{-1}$$

and

$$\begin{aligned} A_{n,1}(1, 0; x) &= \int_x^{1/n} \frac{\left(\frac{1}{n} - u\right)^2}{u^{3/2}} u \, du \, s_{n,1}(x) \\ &= e^{-nx} \left(-2(1 - nx)^2 \frac{x^{3/2}}{n} + \frac{8}{3} (1 - nx)x^{5/2} - \frac{8}{3} nx \int_x^{1/n} u^{3/2} \, du \right) \\ &\leq \frac{c}{n^{3/2}} w(-1, 0; x)^{-1}. \end{aligned}$$

In the other two cases (2.25) follows from the inequality

$$A_{n,k}(\alpha, r; x) \leq c A_{n,k}(\alpha, 0; x)(1 + x)^r, \quad x \in [0, 1],$$

and the estimates we have already established. \square

The last two propositions contain Bernstein-type inequalities.

Proposition 2.4. *Let $w = w(\gamma_0, \gamma_\infty)$ be given by (1.1) with $\gamma_0 \in [-1, 0]$ and $\gamma_\infty \in \mathbb{R}$. Then*

$$\|w\varphi D^2(S_n f)\| \leq c_4 n \|wf\| \quad \forall f \in C(w)(0, \infty) \quad \forall n \geq 1.$$

Proof. Analogously to the assertions we proved so far we shall show that

$$(2.26) \quad |\varphi(x) D^2(S_n f)(x)| \leq c n \|wf\| w(x)^{-1}.$$

To this end, we shall consider the cases $nx \leq 1$ and $nx \geq 1$ separately.

For $nx \leq 1$ we use the representation

$$(2.27) \quad D^2 S_n f(x) = n^2 S_n(\Delta_{1/n}^2 f)(x).$$

It follows from

$$(2.28) \quad D S_n f(x) = n S_n(\Delta_{1/n} f)(x),$$

which is established straightforwardly. By (2.27), we have for $nx \leq 1$

$$(2.29) \quad \begin{aligned} &|\varphi(x) D^2 S_n f(x)| \\ &\leq n^2 x \sum_{k \geq 0} \left(w \left(\frac{k+2}{n} \right)^{-1} + 2w \left(\frac{k+1}{n} \right)^{-1} + w \left(\frac{k}{n} \right)^{-1} \right) s_{n,k}(x) \|wf\|. \end{aligned}$$

But since $nx \leq 1$ and $\gamma_0 \in [-1, 0]$

$$\left(\frac{k+i}{n} \right)^{-\gamma_0} \leq \left(\frac{k}{n} \right)^{-\gamma_0} + c n^{\gamma_0} \leq \left(\frac{k}{n} \right)^{-\gamma_0} + \frac{c}{n} x^{-1-\gamma_0}, \quad i = 1, 2.$$

Consequently,

$$(2.30) \quad w \left(\frac{k+i}{n} \right)^{-1} \leq c w \left(\frac{k}{n} \right)^{-1} + \frac{c}{n} x^{-1-\gamma_0} \left(1 + \frac{k}{n} \right)^{\gamma_0 - \gamma_\infty}, \quad i = 1, 2.$$

Relations (2.29), (2.30) and (2.6) imply (2.26) for $nx \leq 1$.

Let now $nx \geq 1$. In this case we use that

$$(2.31) \quad s'_{n,k}(x) = \frac{n}{x} \left(\frac{k}{n} - x \right) s_{n,k}(x).$$

Therefore

$$D S_n f(x) = \frac{n}{x} S_n((\circ - x)f(\circ))(x);$$

hence

$$D^2 S_n f(x) = \frac{n^2}{x^2} S_n((\circ - x)^2 f(\circ))(x) - \frac{n}{x^2} S_n((\circ - x) f(\circ))(x).$$

Consequently,

$$(2.32) \quad |\varphi(x) D^2 S_n f(x)| \leq \frac{n}{x} [n S_n((\circ - x)^2 w(\circ)^{-1})(x) + S_n(|\circ - x| w(\circ)^{-1})(x)] \|w f\|.$$

For the first summand on the right above we have by Hölder's inequality the estimate (cf. (2.7))

$$(2.33) \quad \begin{aligned} S_n((\circ - x)^2 w(\circ)^{-1})(x) &\leq S_n((\circ - x)^2 w(1, r + 1; \circ))(x)^{\gamma_0(\gamma_\infty - \gamma_0)/r} \\ &\quad \times S_n((\circ - x)^2 w(1, 0; \circ))(x)^{-\gamma_0(1 + (\gamma_\infty - \gamma_0)/r)} \\ &\quad \times S_n((\circ - x)^2 w(0, r; \circ))(x)^{(1 + \gamma_0)(\gamma_0 - \gamma_\infty)/r} \\ &\quad \times S_n((\circ - x)^2 w(0, 0; \circ))(x)^{(1 + \gamma_0)(1 + (\gamma_\infty - \gamma_0)/r)}. \end{aligned}$$

To estimate $S_n((\circ - x)^2 w(1, 0; \circ))(x)$ we observe that by (2.3)

$$(2.34) \quad \begin{aligned} S_n((\circ - x)^2 w(1, 0; \circ))(x) &= x S_n((\circ - x)^2)(x) + S_n((\circ - x)^3)(x) \\ &= \frac{x^2}{n} + \frac{x}{n^2} \leq \frac{2x^2}{n}. \end{aligned}$$

Similarly, we have

$$(2.35) \quad \begin{aligned} S_n((\circ - x)^2 w(1, r + 1; \circ))(x) &\leq c x \sum_{k \geq 0} \left(\frac{k+1}{n} - x \right)^2 \left(1 + \frac{k}{n} \right)^r s_{n,k}(x) \\ &= c x \left[\sum_{k \geq 0} \left(\frac{k}{n} - x \right)^2 \left(1 + \frac{k}{n} \right)^r s_{n,k}(x) + \frac{2}{n} \sum_{k \geq 0} \left(\frac{k}{n} - x \right) \left(1 + \frac{k}{n} \right)^r s_{n,k}(x) \right. \\ &\quad \left. + \frac{1}{n^2} \sum_{k \geq 0} \left(1 + \frac{k}{n} \right)^r s_{n,k}(x) \right] \\ &\leq c \left(\frac{x^2}{n} + \frac{x}{n^2} \right) (1+x)^r \leq \frac{c x^2 (1+x)^r}{n}, \end{aligned}$$

as at the last but one step we have used (2.8), (2.9) and (2.17).

Now, (2.33), (2.3), (2.17), (2.34) and (2.35) yield for $nx \geq 1$

$$(2.36) \quad S_n((\circ - x)^2 w(\circ)^{-1})(x) \leq \frac{c x}{n} w(x)^{-1}.$$

For the second summand on the right of (2.32) we have by the Cauchy inequality, (2.36) and (2.6)

$$\begin{aligned} S_n(|\circ - x| w(\circ)^{-1})(x) &\leq \sqrt{S_n((\circ - x)^2 w(\circ)^{-1})(x)} \sqrt{S_n(w^{-1})(x)} \\ &\leq c \sqrt{\frac{x}{n} w(x)^{-1}} \leq c x w(x)^{-1}, \end{aligned}$$

as $nx \geq 1$.

The inequalities (2.32), (2.36) and the last estimate above imply (2.26) for $nx \geq 1$. This completes the proof of the proposition. \square

The next proposition is the analogue of a result of Knoop and Zhou concerning the Bernstein operator [14, Theorem 2.1]. We prove it following their argument. Perhaps it is worth noting that it fits even more naturally and is easier to apply for the Szász-Mirakjan operator than for the Bernstein operator, for which it was originally developed.

Proposition 2.5. *Let $w = w(\gamma_0, \gamma_\infty)$ be given by (1.1) with $\gamma_0 \in [-1, 0]$ and $\gamma_\infty \in \mathbb{R}$. Let $r \in \mathbb{Z}$ be such that $r/(\gamma_0 - \gamma_\infty) > 1$ if $\gamma_0 \neq \gamma_\infty$, and let $r = 1$ if $\gamma_0 = \gamma_\infty$. We set $\gamma = (1 + \gamma_0)(\gamma_0 - \gamma_\infty)/r$. Let also $m \in \mathbb{N}$ as $m \geq 2$. Then*

$$\|w\varphi^{3/2}D^3(S_n^m g)\| \leq c_5 \sqrt{m^{\gamma-1}(\log m)^{1+\gamma_0}} \sqrt{n} \|w\varphi D^2 g\|$$

$$\forall g \in W^2(w\varphi)(0, \infty) \quad \forall n \geq m^2.$$

The value of the constant c_5 depends only on r .

Proof. For $g \in W^2(w\varphi)(0, \infty)$ the formulae (2.27) and (2.28) give

$$(2.37) \quad DS_n g(x) = \sum_{k \geq 0} n \int_0^{1/n} Dg \left(\frac{k}{n} + u \right) du s_{n,k}(x)$$

and

$$(2.38) \quad D^2 S_n g(x) = \sum_{k \geq 0} n^2 \int_0^{1/n} \int_0^{1/n} D^2 g \left(\frac{k}{n} + u + v \right) du dv s_{n,k}(x).$$

Iterating the latter we arrive at

$$(2.39) \quad D^3 S_n^m g(x) = \sum_{\substack{k_j \geq 0 \\ j=1, \dots, m}} n^2 \int_0^{1/n} \int_0^{1/n} D^2 g \left(\frac{k_1}{n} + u + v \right) du dv \mathcal{S}_{n, \bar{k}} s_{n, k_m}(x),$$

where we have set $\bar{k} = (k_1, \dots, k_m)$,

$$\mathcal{S}_{n, \bar{k}} = \prod_{j=1}^{m-1} s_{n, 2, k_j} \left(\frac{k_{j+1}}{n} \right)$$

and

$$s_{n, i, k}(x) = n^i \int_0^{1/n} \cdots \int_0^{1/n} s_{n, k}(x + t_1 + \cdots + t_i) dt_1 \cdots dt_i.$$

Next, just as in [14, pp. 319-320], using (2.37) and (2.39), we arrive at the following $m-1$ representations of $D^3 S_n^m g$

$$(2.40) \quad D^3 S_n^m g(x) = \sum_{\substack{k_j \geq 0 \\ j=1, \dots, m}} n^2 \int_0^{1/n} \int_0^{1/n} D^2 g \left(\frac{k_1}{n} + u + v \right) du dv \mathcal{S}_{n, \bar{k}} Q_{n, j, \bar{k}} s_{n, k_m}(x), \quad j = 1, \dots, m-1,$$

where

$$Q_{n, j, \bar{k}} = \ell_{n, k_j}^* \left(\frac{k_{j+1}}{n} \right) \ell_{n, k_{j+1}} \left(\frac{k_{j+2}}{n} \right) \cdots \ell_{n, k_{m-1}} \left(\frac{k_m}{n} \right), \quad j = 1, \dots, m-2,$$

$$Q_{n, m-1, \bar{k}} = \ell_{n, k_{m-1}}^* \left(\frac{k_m}{n} \right)$$

and

$$\ell_{n,k}^*(x) = \frac{n \int_0^{1/n} s'_{n,2,k}(x+t) dt}{s_{n,2,k}(x)}, \quad \ell_{n,k}(x) = \frac{s_{n,3,k}(x)}{s_{n,2,k}(x)}.$$

Summing the relations in (2.40) we get

$$(2.41) \quad D^3 S_n^m g(x) = \frac{1}{m-1} \sum_{\substack{k_j \geq 0 \\ j=1, \dots, m}} n^2 \int_0^{1/n} \int_0^{1/n} D^2 g \left(\frac{k_1}{n} + u + v \right) du dv \mathcal{S}_{n,\bar{k}} Q_{n,\bar{k}} s_{n,k_m}(x)$$

with

$$Q_{n,\bar{k}} = \sum_{j=1}^{m-1} Q_{n,j,\bar{k}}.$$

Further, let us observe that

$$(2.42) \quad \left| n^2 \int_0^{1/n} \int_0^{1/n} D^2 g \left(\frac{k}{n} + u + v \right) du dv \right| \leq n^2 \int_0^{1/n} \int_0^{1/n} w \left(\frac{k}{n} + u + v \right)^{-1} \varphi \left(\frac{k}{n} + u + v \right)^{-1} du dv \|w\varphi D^2 g\|$$

and

$$(2.43) \quad n^2 \int_0^{1/n} \int_0^{1/n} w \left(\frac{k}{n} + u + v \right)^{-1} \varphi \left(\frac{k}{n} + u + v \right)^{-1} du dv \leq c w \left(\frac{k+1}{n} \right)^{-1} \varphi \left(\frac{k+1}{n} \right)^{-1}.$$

The first of these estimates is obvious. To verify the second one we proceed as follows:

$$\begin{aligned} & n^2 \int_0^{1/n} \int_0^{1/n} w \left(\frac{k}{n} + u + v \right)^{-1} \varphi \left(\frac{k}{n} + u + v \right)^{-1} du dv \\ &= n^2 \int_{-1/n}^{1/n} \left(\frac{1}{n} - |u| \right) w \left(\frac{k+1}{n} + u \right)^{-1} \varphi \left(\frac{k+1}{n} + u \right)^{-1} du \\ &\leq c \left(1 + \frac{k+1}{n} \right)^{\gamma_0 - \gamma_\infty} n^2 \int_{-1/n}^{1/n} \left(\frac{1}{n} - |u| \right) \left(\frac{k+1}{n} + u \right)^{-\gamma_0 - 1} du \\ &\leq c w \left(\frac{k+1}{n} \right)^{-1} \varphi \left(\frac{k+1}{n} \right)^{-1}, \end{aligned}$$

as at the last estimate we have used that the function $(1/n - |u|)[(k+1)/n + u]^{-\gamma_0 - 1}$ is increasing on $[-1/n, 0]$ and decreasing on $[0, 1/n]$. We derive from (2.41) by the Cauchy inequality and (2.42)-(2.43) the estimate

$$|D^3 S_n^m g(x)| \leq \frac{1}{m-1} \sqrt{\sum_{\substack{k_j \geq 0 \\ j=1, \dots, m}} \mathcal{S}_{n,\bar{k}} Q_{n,\bar{k}}^2 s_{n,k_m}(x)}$$

$$\begin{aligned}
 & \times \sqrt{\sum_{\substack{k_j \geq 0 \\ j=1, \dots, m}} \left| n^2 \int_0^{1/n} \int_0^{1/n} D^2 g \left(\frac{k_1}{n} + u + v \right) du dv \right|^2} \mathcal{S}_{n, \bar{k}} s_{n, k_m}(x) \\
 & \leq \frac{c}{m} \sqrt{\sum_{\substack{k_j \geq 0 \\ j=1, \dots, m}} \mathcal{S}_{n, \bar{k}} Q_{n, \bar{k}}^2 s_{n, k_m}(x)} \\
 & \times \sqrt{\sum_{\substack{k_j \geq 0 \\ j=1, \dots, m}} w \left(\frac{k_1 + 1}{n} \right)^{-2} \varphi \left(\frac{k_1 + 1}{n} \right)^{-2} \mathcal{S}_{n, \bar{k}} s_{n, k_m}(x) \|w\varphi D^2 g\|}.
 \end{aligned}$$

Thus we can finish the proof of the proposition if we show that

$$(2.44) \quad \sum_{\substack{k_j \geq 0 \\ j=1, \dots, m}} \mathcal{S}_{n, \bar{k}} Q_{n, \bar{k}}^2 s_{n, k_m}(x) \leq c n m \varphi(x)^{-1}$$

and

$$(2.45) \quad \sum_{\substack{k_j \geq 0 \\ j=1, \dots, m}} w \left(\frac{k_1 + 1}{n} \right)^{-2} \varphi \left(\frac{k_1 + 1}{n} \right)^{-2} \mathcal{S}_{n, \bar{k}} s_{n, k_m}(x) \leq c w(x)^{-2} \varphi(x)^{-2} m^\gamma (\log m)^{1+\gamma_0}, \quad n \geq m^2.$$

In order to simplify these two assertions we take into account that $s_{n, i, k}(x) = s_{1, i, k}(nx)$. Consequently,

$$\begin{aligned}
 \mathcal{S}_{n, \bar{k}} &= \mathcal{S}_{1, \bar{k}}, \\
 \ell_{n, k}^*(x) &= n \ell_{1, k}^*(nx), \quad \ell_{n, k}(x) = \ell_{1, k}(nx), \\
 Q_{n, j, \bar{k}} &= n Q_{1, j, \bar{k}}.
 \end{aligned}$$

Also, we have (cf. [14, pp. 327-328])

$$\sum_{\substack{k_j \geq 0 \\ j=1, \dots, m}} \mathcal{S}_{n, \bar{k}} Q_{n, j', \bar{k}} Q_{n, j'', \bar{k}} s_{n, k_m}(x) \equiv 0, \quad j' \neq j''.$$

Hence (2.44) and (2.45) reduce respectively to

$$\sum_{\substack{k_j \geq 0 \\ j=1, \dots, m}} \mathcal{S}_{1, \bar{k}} Q_{1, j, \bar{k}}^2 s_{1, k_m}(x) \leq c \varphi(x)^{-1}, \quad j = 1, \dots, m-1,$$

and

$$(2.46) \quad \sum_{\substack{k_j \geq 0 \\ j=1, \dots, m}} (1 + k_1)^{-2(1+\gamma_0)} (n + k_1)^{2(\gamma_0 - \gamma_\infty)} \mathcal{S}_{1, \bar{k}} s_{1, k_m}(x) \leq c x^{-2(1+\gamma_0)} (n + x)^{2(\gamma_0 - \gamma_\infty)} m^\gamma (\log m)^{1+\gamma_0}, \quad n \geq m^2.$$

The first estimate is verified in Lemma 2.6 below. As for the second we use Hölder's inequality as in the proof of the previous propositions to get

$$\begin{aligned}
& \sum_{\substack{k_j \geq 0 \\ j=1, \dots, m}} (1+k_1)^{-2(1+\gamma_0)} (n+k_1)^{2(\gamma_0-\gamma_\infty)} \mathcal{S}_{1, \bar{k}} s_{1, k_m}(x) \\
& \leq \left\{ \sum_{\substack{k_j \geq 0 \\ j=1, \dots, m}} (1+k_1)^{-4} \mathcal{S}_{1, \bar{k}} s_{1, k_m}(x) \right\}^{(1+\gamma_0)(\gamma_0-\gamma_\infty)/(2r)} \\
& \quad \times \left\{ \sum_{\substack{k_j \geq 0 \\ j=1, \dots, m}} (1+k_1)^{-2} \mathcal{S}_{1, \bar{k}} s_{1, k_m}(x) \right\}^{(1+\gamma_0)(1+(\gamma_\infty-\gamma_0)/r)} \\
& \quad \times \left\{ \sum_{\substack{k_j \geq 0 \\ j=1, \dots, m}} (n+k_1)^{2r} \mathcal{S}_{1, \bar{k}} s_{1, k_m}(x) \right\}^{\gamma_0(\gamma_\infty-\gamma_0)/r} \\
& \quad \times \left\{ \sum_{\substack{k_j \geq 0 \\ j=1, \dots, m}} (n+k_1)^{4r} \mathcal{S}_{1, \bar{k}} s_{1, k_m}(x) \right\}^{(1+\gamma_0)(\gamma_0-\gamma_\infty)/(2r)} \\
& \quad \times \left\{ \sum_{\substack{k_j \geq 0 \\ j=1, \dots, m}} \mathcal{S}_{1, \bar{k}} s_{1, k_m}(x) \right\}^{-\gamma_0(1+(\gamma_\infty-\gamma_0)/r)}.
\end{aligned}$$

Now, (2.46) follows from Lemma 2.7 and the fact that the last term above is equal to 1. \square

Now we shall establish the two lemmas used in the proof above.

Lemma 2.6. *Let $m \in \mathbb{N}$ as $m \geq 2$. Then there hold*

$$\sum_{\substack{k_j \geq 0 \\ j=1, \dots, m}} \mathcal{S}_{1, \bar{k}} Q_{1, j, \bar{k}}^2 s_{1, k_m}(x) \leq c \varphi(x)^{-1}, \quad x > 0,$$

for $j = 1, \dots, m-1$. The value of the constant c does not depend on x or m .

Proof. We follow the argument of Knoop and Zhou [14]. We start with the equality

$$\begin{aligned}
(2.47) \quad & \sum_{\substack{k_j \geq 0 \\ j=1, \dots, m}} \mathcal{S}_{1, \bar{k}} Q_{1, j, \bar{k}}^2 s_{1, k_m}(x) = \sum_{k_m \geq 0} s_{1, k_m}(x) \sum_{k_{m-1} \geq 0} s_{1, 2, k_{m-1}}(k_m) \ell_{1, k_{m-1}}(k_m)^2 \\
& \quad \cdots \sum_{k_{j+1} \geq 0} s_{1, 2, k_{j+1}}(k_{j+2}) \ell_{1, k_{j+1}}(k_{j+2})^2 \sum_{k_j \geq 0} s_{1, 2, k_j}(k_{j+1}) \ell_{1, k_j}^*(k_{j+1})^2.
\end{aligned}$$

Next, we establish that

$$(2.48) \quad \sum_{k \geq 0} s_{1,2,k}(i) \ell_{1,k}^*(i)^2 \leq \frac{c}{1+i}, \quad i \in \mathbb{N}_0,$$

and

$$(2.49) \quad \sum_{k \geq 0} s_{1,2,k}(i) \ell_{1,k}(i)^2 \frac{1}{1+k} \leq \frac{1}{1+i}, \quad i \in \mathbb{N}_0.$$

Then the lemma follows from (2.47)-(2.49) and (2.8) (with $r = -1$).

To prove (2.48) for $i = 0$ we use that $s'_{n,k}(x) = n(s_{n,k-1}(x) - s_{n,k}(x))$, $k > 0$, to derive the estimate

$$\left| \int_0^1 \int_0^1 \int_0^1 s'_{1,k}(t_1 + t_2 + t_3) dt_1 dt_2 dt_3 \right| \leq \frac{3^k(k+1)}{k!}.$$

Since, also, we have

$$\int_0^1 \int_0^1 s_{1,k}(t_1 + t_2) dt_1 dt_2 \geq \frac{1}{e^2 k!} \int_0^1 t^k dt = \frac{1}{e^2(k+1)!},$$

we arrive at

$$\begin{aligned} s_{1,2,k}(0) \ell_{1,k}^*(0)^2 &= \frac{\left(\int_0^1 \int_0^1 \int_0^1 s'_{1,k}(t_1 + t_2 + t_3) dt_1 dt_2 dt_3 \right)^2}{\int_0^1 \int_0^1 s_{1,k}(t_1 + t_2) dt_1 dt_2} \\ &\leq \frac{e^2 3^{2k} (k+1)^3}{k!}; \end{aligned}$$

hence (2.48) follows for $i = 0$.

To prove (2.48) for $i \geq 1$ we apply the formula (2.31) and the Cauchy inequality to get

(2.50)

$$\begin{aligned} s_{1,2,k}(i) \ell_{1,k}^*(i)^2 &= \frac{\left(\int_0^1 \int_0^1 \int_0^1 s'_{1,k}(i + t_1 + t_2 + t_3) dt_1 dt_2 dt_3 \right)^2}{\int_0^1 \int_0^1 s_{1,k}(i + t_1 + t_2) dt_1 dt_2} \\ &\leq \int_0^1 \int_0^1 \int_0^1 \left(\frac{k - (i + t_1 + t_2 + t_3)}{i + t_1 + t_2 + t_3} \right)^2 \frac{s_{1,k}(i + t_1 + t_2 + t_3)^2}{s_{1,k}(i + t_1 + t_2)} dt_1 dt_2 dt_3 \\ &\leq \frac{c}{i^2} ([k - (i + 5)]^2 + 1) s_{1,k}(i + 5) \end{aligned}$$

since for $t_1, t_2, t_3 \in [0, 1]$

$$\left(\frac{k - (i + t_1 + t_2 + t_3)}{i + t_1 + t_2 + t_3} \right)^2 \leq \frac{c}{i^2} ([k - (i + 5)]^2 + 1)$$

and

$$\frac{(i + t_1 + t_2 + t_3)^2}{i + t_1 + t_2} \leq i + 5, \quad i \geq 1.$$

Now, (2.48) follows for $i \geq 1$ from (2.50), (2.2) and (2.3).

The inequality (2.49) is derived by similar though more precise estimates. First, let $i = 0$. We observe that

$$\begin{aligned}
\int_0^1 \int_0^1 s_{1,k}(t_1 + t_2) dt_1 dt_2 &= \frac{1}{k!} \int_0^1 \int_0^1 e^{-t_1-t_2} (t_1 + t_2)^k dt_1 dt_2 \\
&= \frac{1}{k!} \sum_{l=0}^k \binom{k}{l} \int_0^1 e^{-t_1} t_1^l dt_1 \int_0^1 e^{-t_2} t_2^{k-l} dt_2 \\
&\geq \frac{1}{k!} \sum_{l=0}^k \binom{k}{l} \int_0^1 e^{-t_1} t_1^l dt_1 \frac{1}{e^{(k-l+1)}} \\
&\geq \frac{1}{e^{(k+1)}} \frac{1}{k!} \int_0^1 e^{-t_1} \sum_{l=0}^k \binom{k}{l} t_1^l dt_1 \\
&= \frac{1}{k+1} \int_0^1 s_{1,k}(1+t_1) dt_1.
\end{aligned}$$

Then by the Cauchy inequality we get

$$\begin{aligned}
s_{1,2,k}(0) \ell_{1,k}(0)^2 \frac{1}{1+k} &= \frac{1}{1+k} \frac{\left(\int_0^1 \int_0^1 \int_0^1 s_{1,k}(t_1 + t_2 + t_3) dt_1 dt_2 dt_3 \right)^2}{\int_0^1 \int_0^1 s_{1,k}(t_1 + t_2) dt_1 dt_2} \\
&\leq \int_0^1 \int_0^1 \int_0^1 \frac{s_{1,k}(t_1 + t_2 + t_3)^2}{s_{1,k}(1+t_1)} dt_1 dt_2 dt_3 \\
&= \int_0^1 \int_0^1 \int_0^1 e^{1-t_1-2t_2-2t_3} \frac{1}{k!} \left[\frac{(t_1 + t_2 + t_3)^2}{1+t_1} \right]^k dt_1 dt_2 dt_3.
\end{aligned}$$

Now, to estimate the sum on the left hand-side of (2.49) for $i = 0$ we calculate separately its term for $k = 0$ whereas for the terms with $k \geq 1$ we apply the inequality above. Thus we get

$$\begin{aligned}
&\sum_{k \geq 0} s_{1,2,k}(0) \ell_{1,k}(0)^2 \frac{1}{1+k} \\
&\leq (1 - e^{-1})^4 + \int_0^1 \int_0^1 \int_0^1 \left(e^{\frac{(1-t_2-t_3)^2}{1+t_1}} - e^{1-t_1-2t_2-2t_3} \right) dt_1 dt_2 dt_3 \\
&= \int_0^1 \int_0^1 \int_0^1 e^{\frac{(1-t_2-t_3)^2}{1+t_1}} dt_1 dt_2 dt_3 - \frac{e}{4} (1 - e^{-1})(1 - e^{-2})^2 + (1 - e^{-1})^4,
\end{aligned}$$

hence, in view of [14, (4.9)], (2.49) follows for $i = 0$.

For $i \geq 1$ we have by the Cauchy inequality

$$\begin{aligned} s_{1,2,k}(i) \ell_{1,k}(i)^2 \frac{1}{1+k} &= \frac{1}{1+k} \frac{\left(\int_0^1 \int_0^1 \int_0^1 s_{1,k}(i+t_1+t_2+t_3) dt_1 dt_2 dt_3 \right)^2}{\int_0^1 \int_0^1 s_{1,k}(i+t_1+t_2) dt_1 dt_2} \\ &\leq \frac{1}{1+k} \int_0^1 \int_0^1 \int_0^1 \frac{s_{1,k}(i+t_1+t_2+t_3)^2}{s_{1,k}(i+t_1+t_2)} dt_1 dt_2 dt_3 \\ &= \int_0^1 \int_0^1 \int_0^1 \frac{i+t_1+t_2}{(i+t_1+t_2+t_3)^2} e^{\frac{t_3^2}{i+t_1+t_2}} s_{1,k+1} \left(\frac{(i+t_1+t_2+t_3)^2}{i+t_1+t_2} \right) dt_1 dt_2 dt_3. \end{aligned}$$

Summing on $k \geq 0$ we arrive at the estimate

$$\sum_{k \geq 0} s_{1,2,k}(i) \ell_{1,k}(i)^2 \frac{1}{1+k} \leq \int_0^1 \int_0^1 \int_0^1 \frac{i+t_1+t_2}{(i+t_1+t_2+t_3)^2} e^{\frac{t_3^2}{i+t_1+t_2}} dt_1 dt_2 dt_3.$$

Now, (2.49) follows for $i \geq 1$ from [14, (4.10)]. \square

Lemma 2.7. *Let $m \in \mathbb{N}$ as $m \geq 2$.*

(a) *For $i \in \mathbb{Z}_-$ we have*

$$\sum_{\substack{k_j \geq 0 \\ j=1, \dots, m}} (1+k_1)^i \mathcal{S}_{1, \bar{k}} s_{1, k_m}(x) \leq c x^i m^{-i-2} \log m, \quad x > 0.$$

(b) *For $i \in \mathbb{Z}_-$ we have*

$$\sum_{\substack{k_j \geq 0 \\ j=1, \dots, m}} (n+k_1)^i \mathcal{S}_{1, \bar{k}} s_{1, k_m}(x) \leq c(n+x)^i, \quad x \geq 0, \quad n \geq m^2.$$

(c) *For $i \in \mathbb{N}_0$ we have*

$$\sum_{\substack{k_j \geq 0 \\ j=1, \dots, m}} (n+k_1)^i \mathcal{S}_{1, \bar{k}} s_{1, k_m}(x) \leq c(n+x)^i, \quad x \geq 0, \quad n \geq m.$$

The value of the constant c depends only on i .

Proof. We again follow the argument of Knoop and Zhou [14]. For $i \leq -1$ we set $\ell = -i$. We shall use the representation

$$(n+k)^i = \int_0^1 \cdots \int_0^1 (\tau_1 \cdots \tau_\ell)^{n+k-1} d\tau_1 \cdots d\tau_\ell$$

and the formula

$$\sum_{k \geq 0} a^k s_{1,2,k}(x) = e^{-(1-a)x} \left(\frac{1 - e^{-(1-a)}}{1-a} \right)^2.$$

The latter is checked directly by means of the argument

$$\begin{aligned} \sum_{k \geq 0} a^k s_{1,2,k}(x) &= \int_0^1 \int_0^1 e^{-x-t_1-t_2} \sum_{k \geq 0} \frac{[a(x+t_1+t_2)]^k}{k!} dt_1 dt_2 \\ &= \int_0^1 \int_0^1 e^{(a-1)(x+t_1+t_2)} dt_1 dt_2 \end{aligned}$$

$$= e^{(a-1)x} \left(\int_0^1 e^{(a-1)t} dt \right)^2.$$

Now, just as in [14, pp. 322-323], we arrive at

$$\sum_{\substack{k_j \geq 0 \\ j=1, \dots, m}} (n + k_1)^i \mathcal{S}_{1, \bar{k}} s_{1, k_m}(x) \\ = \int_0^1 \dots \int_0^1 (\tau_1 \dots \tau_\ell)^{n-1} \frac{F_{m-1}(\tau_1 \dots \tau_\ell)^2}{F_0(\tau_1 \dots \tau_\ell)^2} e^{-F_{m-1}(\tau_1 \dots \tau_\ell)x} d\tau_1 \dots d\tau_\ell,$$

where

$$F_0(\tau) = 1 - \tau, \quad F_\rho(\tau) = 1 - e^{-F_{\rho-1}(\tau)}, \quad \rho = 1, 2, \dots$$

Let $\mathcal{D} \subset [0, 1]^\ell$ be a rectangle with at least one side of the form $[0, 1/2]$. Then, clearly,

$$(2.51) \quad F_0(\tau_1 \dots \tau_\ell) \geq 1/2, \quad (\tau_1, \dots, \tau_\ell) \in \mathcal{D}.$$

It can be shown by induction that

$$(2.52) \quad \frac{1 - \tau}{m} \leq F_{m-1}(\tau) \leq 1, \quad 0 \leq \tau \leq 1.$$

Also, since $y^\ell e^{-y} \leq c$, $y \geq 0$, then

$$(2.53) \quad F_{m-1}(\tau)^\ell e^{-F_{m-1}(\tau)x} \leq c x^i, \quad \tau \geq 0, \quad x > 0.$$

Now, (2.51), the left inequality in (2.52) and (2.53) imply

$$(2.54) \quad \int_{\mathcal{D}} \frac{F_{m-1}(\tau_1 \dots \tau_\ell)^2}{F_0(\tau_1 \dots \tau_\ell)^2} e^{-F_{m-1}(\tau_1 \dots \tau_\ell)x} d\tau_1 \dots d\tau_\ell \leq c m^{-i-2} x^i.$$

Similarly, (2.51) and the right inequality in (2.52) directly imply the relation

$$(2.55) \quad \int_{\mathcal{D}} (\tau_1 \dots \tau_\ell)^{n-1} \frac{F_{m-1}(\tau_1 \dots \tau_\ell)^2}{F_0(\tau_1 \dots \tau_\ell)^2} e^{-F_{m-1}(\tau_1 \dots \tau_\ell)x} d\tau_1 \dots d\tau_\ell \leq c n^i, \quad n \geq 1.$$

For $n \geq m$ (2.51), the left inequality in (2.52) and (2.53) yield

$$(2.56) \quad \int_{\mathcal{D}} (\tau_1 \dots \tau_\ell)^{n-1} \frac{F_{m-1}(\tau_1 \dots \tau_\ell)^2}{F_0(\tau_1 \dots \tau_\ell)^2} e^{-F_{m-1}(\tau_1 \dots \tau_\ell)x} d\tau_1 \dots d\tau_\ell \\ \leq c m^{\ell-2} (nx)^i \leq c x^i, \quad n \geq m.$$

Now, by means of (2.54) we derive

$$\int_0^1 \dots \int_0^1 \frac{F_{m-1}(\tau_1 \dots \tau_\ell)^2}{F_0(\tau_1 \dots \tau_\ell)^2} e^{-F_{m-1}(\tau_1 \dots \tau_\ell)x} d\tau_1 \dots d\tau_\ell \\ \leq \int_{1/2}^1 \dots \int_{1/2}^1 \frac{F_{m-1}(\tau_1 \dots \tau_\ell)^2}{F_0(\tau_1 \dots \tau_\ell)^2} e^{-F_{m-1}(\tau_1 \dots \tau_\ell)x} d\tau_1 \dots d\tau_\ell + c m^{-i-2} x^i;$$

and similarly by (2.55)-(2.56) we get

$$\int_0^1 \dots \int_0^1 (\tau_1 \dots \tau_\ell)^{n-1} \frac{F_{m-1}(\tau_1 \dots \tau_\ell)^2}{F_0(\tau_1 \dots \tau_\ell)^2} e^{-F_{m-1}(\tau_1 \dots \tau_\ell)x} d\tau_1 \dots d\tau_\ell \\ \leq \int_{1/2}^1 \dots \int_{1/2}^1 (\tau_1 \dots \tau_\ell)^{n-1} \frac{F_{m-1}(\tau_1 \dots \tau_\ell)^2}{F_0(\tau_1 \dots \tau_\ell)^2} e^{-F_{m-1}(\tau_1 \dots \tau_\ell)x} d\tau_1 \dots d\tau_\ell \\ + c (n+x)^i, \quad n \geq m.$$

So, to complete the proof of (a) and (b) it is enough to show respectively that that

$$(2.57) \quad \int_{1/2}^1 \cdots \int_{1/2}^1 \frac{F_{m-1}(\tau_1 \cdots \tau_\ell)^2}{F_0(\tau_1 \cdots \tau_\ell)^2} e^{-F_{m-1}(\tau_1 \cdots \tau_\ell)x} d\tau_1 \cdots d\tau_\ell \leq c x^i m^{-i-2} \log m,$$

and

$$(2.58) \quad \int_{1/2}^1 \cdots \int_{1/2}^1 (\tau_1 \cdots \tau_\ell)^{n-1} \frac{F_{m-1}(\tau_1 \cdots \tau_\ell)^2}{F_0(\tau_1 \cdots \tau_\ell)^2} e^{-F_{m-1}(\tau_1 \cdots \tau_\ell)x} d\tau_1 \cdots d\tau_\ell \leq c(n+x)^i, \quad n \geq m^2.$$

We set

$$F_n(\tau, x) = \tau^{n-1} \frac{F_{m-1}(\tau)^2}{F_0(\tau)^2} e^{-F_{m-1}(\tau)x}.$$

In the integral in (2.58) we make the change of the variables, defined by the formulae $\sigma_\rho = \tau_1 \cdots \tau_\rho$, $\rho = 1, \dots, \ell$, and arrange the order of integration from σ_1 to σ_ℓ to get the estimate

$$(2.59) \quad \begin{aligned} & \int_{1/2}^1 \cdots \int_{1/2}^1 F_n(\tau_1 \cdots \tau_\ell, x) d\tau_1 \cdots d\tau_\ell \\ & \leq \int_{2^i}^1 \left[F_n(\sigma_\ell, x) \int_{\sigma_\ell}^1 \left(\frac{1}{\sigma_{\ell-1}} \cdots \left(\frac{1}{\sigma_3} \int_{\sigma_3}^1 \left(\frac{1}{\sigma_2} \int_{\sigma_2}^1 \frac{1}{\sigma_1} d\sigma_1 \right) d\sigma_2 \right) \cdots \right) d\sigma_{\ell-1} \right] d\sigma_\ell \\ & \leq c \int_{2^i}^1 \left[F_n(\sigma_\ell, x) \int_{\sigma_\ell}^1 \left(\cdots \left(\int_{\sigma_3}^1 \left(\int_{\sigma_2}^1 d\sigma_1 \right) d\sigma_2 \right) \cdots \right) d\sigma_{\ell-1} \right] d\sigma_\ell \\ & \leq c \int_{2^i}^1 F_n(\sigma_\ell, x) (1 - \sigma_\ell)^{\ell-1} d\sigma_\ell \\ & \leq c \int_0^1 (1-v)^{n-1} v^{\ell-3} G_{m-1}(v)^2 e^{-G_{m-1}(v)x} dv, \end{aligned}$$

as at the last step we have made the change of the variable $\sigma_\ell = 1 - v$ and set

$$G_0(v) = v, \quad G_\rho(v) = 1 - e^{-G_{\rho-1}(v)}, \quad \rho = 1, 2, \dots$$

Routine considerations show that

$$(2.60) \quad v - \frac{m}{2} v^2 \leq G_{m-1}(v) \leq v, \quad 0 \leq v \leq 1.$$

For $n = 1$ (2.59) reads

$$\begin{aligned} & \int_{1/2}^1 \cdots \int_{1/2}^1 \frac{F_{m-1}(\tau_1 \cdots \tau_\ell)^2}{F_0(\tau_1 \cdots \tau_\ell)^2} e^{-F_{m-1}(\tau_1 \cdots \tau_\ell)x} d\tau_1 \cdots d\tau_\ell \\ & \leq c \int_0^1 v^{\ell-3} G_{m-1}(v)^2 e^{-G_{m-1}(v)x} dv. \end{aligned}$$

We split the integral on the right by means of the intermediate point $1/m$. For the one between 0 and $1/m$ we apply (2.60) to get

$$(2.61) \quad \int_0^{1/m} v^{\ell-3} G_{m-1}(v)^2 e^{-G_{m-1}(v)x} dv \leq c \int_0^1 v^{\ell-1} e^{-vx} dv \leq c x^i,$$

as the last estimate is verified by integration by parts.

For the other integral we again use (2.52) and (2.53) to arrive at

$$(2.62) \quad \int_{1/m}^1 v^{\ell-3} G_{m-1}(v)^2 e^{-G_{m-1}(v)x} dv \leq c m^{\ell-2} x^i \int_{1/m}^1 \frac{dv}{v} = c x^i m^{-i-2} \log m.$$

Now, (2.61)-(2.62) imply (2.57), which completes the proof of assertion (a).

The estimate (2.58) is established in a similar way. Using the right inequality in (2.60), we get

$$(2.63) \quad \int_0^1 (1-v)^{n-1} v^{\ell-3} G_{m-1}(v)^2 e^{-G_{m-1}(v)x} dv \leq \int_0^1 (1-v)^{n-1} v^{\ell-1} dv \\ = \frac{(\ell-1)!(n-1)!}{(\ell+n-1)!} \leq c n^i.$$

As above we get

$$(2.64) \quad \int_0^{1/m} (1-v)^{n-1} v^{\ell-3} G_{m-1}(v)^2 e^{-G_{m-1}(v)x} dv \leq c \int_0^1 v^{\ell-1} e^{-vx} dv \leq c x^i.$$

To estimate the integral between $1/m$ and 1, we first observe that

$$(2.65) \quad m^\ell \left(1 - \frac{1}{m}\right)^{m^2} \leq m^\ell e^{-m} \leq c,$$

which follow from $(1-y)^n \leq e^{-ny}$, $y \in [0, 1]$, and $y^\ell e^{-y} \leq c$, $y \geq 0$. Now, by means of the first inequality in (2.52), (2.53) and (2.65), we get for $n \geq m^2$

$$(2.66) \quad \int_{1/m}^1 (1-v)^{n-1} v^{\ell-3} G_{m-1}(v)^2 e^{-G_{m-1}(v)x} dv \\ \leq c \left(1 - \frac{1}{m}\right)^{n-1} m^{\ell-1} x^i \leq c m^\ell e^{-m} x^i \leq c x^i, \quad n \geq m^2.$$

The estimates (2.59), (2.63), (2.64) and (2.66) imply (2.58). Thus assertion (b) is established.

For the proof of (c) we first observe that for $\theta \geq 0$ there holds

$$\sum_{k \geq 0} (\theta + k)^i s_{1,k}(x) \leq (\theta + x) \sum_{k \geq 0} (\theta + 1 + k)^{i-1} s_{1,k}(x).$$

Iterating it we arrive at

$$(2.67) \quad \sum_{k \geq 0} (\theta + k)^i s_{1,k}(x) \leq (\theta + x)(\theta + 1 + x) \cdots (\theta + i - 1 + x) \leq (\theta + i + x)^i.$$

Similarly,

$$\sum_{k \geq 0} (\theta + k)^i s_{1,2,k}(x) \leq (\theta + 2 + x) \sum_{k \geq 0} (\theta + 1 + k)^{i-1} s_{1,2,k}(x);$$

hence

$$(2.68) \quad \sum_{k \geq 0} (\theta + k)^i s_{1,2,k}(x) \leq (\theta + 2 + x)(\theta + 3 + x) \cdots (\theta + i + 1 + x) \\ \leq (\theta + i + 1 + x)^i.$$

Now, applying (2.68) consecutively with $\theta = n, n + i + 1, \dots, n + (m - 2)(i + 1)$ and then (2.67) with $\theta = n + (m - 1)(i + 1)$ we get

$$\begin{aligned}
 & \sum_{\substack{k_j \geq 0 \\ j=1, \dots, m}} (n + k_1)^i \mathcal{S}_{1, \bar{k}} s_{1, k_m}(x) \\
 &= \sum_{k_m \geq 0} s_{1, k_m}(x) \sum_{k_{m-1} \geq 0} s_{1, 2, k_{m-1}}(k_m) \cdots \sum_{k_1 \geq 0} (n + k_1)^i s_{1, 2, k_1}(k_2) \\
 &\leq \sum_{k_m \geq 0} s_{1, k_m}(x) \sum_{k_{m-1} \geq 0} s_{1, 2, k_{m-1}}(k_m) \cdots \sum_{k_2 \geq 0} (n + i + 1 + k_2)^i s_{1, 2, k_2}(k_3) \\
 &\leq \sum_{k_m \geq 0} [n + (m - 1)(i + 1) + k_m]^i s_{1, k_m}(x) \\
 &\leq [n + (m - 1)(i + 1) + i + x]^i,
 \end{aligned}$$

which, in view of $n \geq m$, implies assertion (c) of the lemma. \square

We are ready to prove our main result.

Proof of Theorem 1.2. It is enough to prove the theorem for $f \in C(w)(0, \infty)$ since $\|w(f - S_n f)\|$ and $K_w(f, n^{-1})$ are invariant under addition of a liner function to f .

The upper error estimate of the theorem follows from Propositions 2.1 and 2.2 via a direct and well-known argument.

To verify the strong converse inequality we apply [3, Theorem 4.1] with $Q_\alpha = S_n$, $\alpha = n^{-1}$, $X = C(w)[0, \infty)$, $Y = W^2(w\varphi)(0, \infty)$, $Z = W^3(w\varphi^{3/2})(0, \infty)$, the differential operator D of [3, Theorem 4.1] is φD^2 , $\Phi(g) = \|w\varphi^{3/2} D^3 g\|$, $M = c_1$, $\lambda(\alpha) = (2n)^{-1}$, $\lambda_1(\alpha) = c_3 n^{-3/2}$, $\ell = 1$, $B = c_4/2$ and m replaced with $m + 1$. Then

$$A = 2c_3 c_5 \sqrt{m^{\gamma-1} (\log m)^{1+\gamma_0}}.$$

We fix $m \in \mathbb{N}$, $m \geq 2$, so large that $A < 1$. Note that $\gamma < 1$. Then [3, Theorem 4.1], Propositions 2.1, 2.3, 2.4 and 2.5 imply the left hand-side inequality of Theorem 1.2 for any $n \geq m^2$. To extend the latter for $1 \leq n \leq m^2 - 1$, we first observe that the property (2.5) of the Szász-Mirakjan operator yields

$$S_n f = [S_{m'}(f_{m'/n})]_{n/m'},$$

where we have set $m' = m^2$.

Consequently,

$$(2.69) \quad \|w(f - S_n f)\| = \|w_{m'/n}(f_{m'/n} - S_{m'}(f_{m'/n}))\| \geq c \|w(f_{m'/n} - S_{m'}(f_{m'/n}))\|$$

with a constant $c > 0$ whose value is independent of f and n because $1 \leq m'/n \leq m'$ and

$$w_\nu(x) \geq \nu^{\min\{\gamma_0, \gamma_\infty\}} w(x), \quad \nu \geq 1.$$

On the other hand,

$$(2.70) \quad K_w(f_\nu, t) = K_{w_{1/\nu}}(f, \nu t) \geq \nu^{-\max\{\gamma_0, \gamma_\infty\}} K_w(f, \nu t), \quad \nu \geq 1,$$

because $(g_\nu)' = \nu(g')_\nu$ and

$$w_{1/\nu}(x) \geq \nu^{-\max\{\gamma_0, \gamma_\infty\}} w(x), \quad \nu \geq 1.$$

Now, combining the relation

$$K_w\left(f, \frac{1}{m'}\right) \leq c \|w(f - S_{m'} f)\| \quad \forall f \in C(w)(0, \infty)$$

with (2.69) and (2.70) with $\nu = m'/n \in [1, m']$ and $t = 1/m'$, we get

$$K_w \left(f, \frac{1}{n} \right) \leq c \|w(f - S_n f)\| \quad \forall f \in C(w)(0, \infty)$$

for $1 \leq n \leq m^2 - 1$ as well. \square

Acknowledgments. We are thankful to Kiril Delev for carefully reading the manuscript and indicating several misprints.

REFERENCES

1. N. T. Amanov, *On the weighted approximation by Szász-Mirakjan operators*, Anal. Math. **18** (1992), 167–184.
2. M. Becker, *Global approximation theorems for Szász-Mirakjan and Baskakov operators in polynomial weight spaces*, Indiana Univ. Math. J. **27** No. 1 (1978), 127–142.
3. Z. Ditzian, K. G. Ivanov, *Strong converse inequalities*, J. D'Analyse Math. **61** (1993), 61–111.
4. Z. Ditzian, V. Totik, *Moduli of smoothness*, Springer Verlag, New York, 1987.
5. B. R. Draganov and K. G. Ivanov, *A new characterization of weighted Peetre K -functionals*, Constr. Approx. **21** (2005), 113–148.
6. B. R. Draganov and K. G. Ivanov, *A new characterization of weighted Peetre K -functionals (II)*, Serdica Math. J. **33** (2007), 59–124.
7. B. R. Draganov and K. G. Ivanov, *Equivalence between K -functionals based on continuous linear transforms*, Serdica Math. J. **33** (2007), 475–494.
8. B. R. Draganov and K. G. Ivanov, *A new characterization of weighted Peetre K -functionals (III)*, manuscript, 2012.
9. B. R. Draganov and K. G. Ivanov, *A new characterization of weighted Peetre K -functionals (IV)*, manuscript, 2012.
10. Z. Finta, *On converse approximation theorems*, J. Math. Anal. Appl. **312** (2005), 159–180.
11. A. Holhoş, *Uniform weighted approximation by positive linear operators*, Stud. Univ. Babeş-Bolyai Math. **56** (2011), No. 3, 135–146.
12. K. G. Ivanov, *A characterization of weighted Peetre K -functionals*, J. Approx. Theory **56** (1989), 185–211.
13. K. G. Ivanov, *Natural weights for uniform approximation by Bernstein polynomials*, manuscript, 2011.
14. H.-B. Knoop and X.-L. Zhou, *The lower estimate for linear positive operators (II)*, Results Math., **25** (1994), 315–330.
15. V. Totik, *Approximation by Bernstein polynomials*, Amer. J. Math., **116** (1994), 995–1018.

DEPARTMENT OF MATHEMATICS AND INFORMATICS, UNIVERSITY OF SOFIA, 5 JAMES BOURCHIER BLVD., 1164 SOFIA, BULGARIA; INSTITUTE OF MATHEMATICS AND INFORMATICS, BULGARIAN ACADEMY OF SCIENCES, BL. 8 ACAD. G. BONCHEV STR., 1113 SOFIA, BULGARIA

E-mail address: `bdraganov@fmi.uni-sofia.bg`

INSTITUTE OF MATHEMATICS AND INFORMATICS, BULGARIAN ACADEMY OF SCIENCES, BL. 8 ACAD. G. BONCHEV STR., 1113 SOFIA, BULGARIA

E-mail address: `kamen@math.bas.bg`