

An improved Jackson inequality for best trigonometric approximation

Borislav R. Draganov*

Abstract

The paper presents an improved Jackson inequality and a corresponding inverse one for best trigonometric approximation in terms of moduli of smoothness that are equivalent to zero on the trigonometric polynomials up to a certain degree. The inequalities are analogous to M.F. Timan's. Relations between such moduli of different order are also considered.

AMS classification: 41A17, 41A27, 42A10

Key words and phrases: best trigonometric approximation, Jackson inequality, Timan's inequality, modulus of smoothness, linear transform.

1 Introduction

Let $L_p(\mathbb{T})$, $1 \leq p \leq \infty$, denote the space of the functions with finite L_p -norm on the circle \mathbb{T} . We can consider $C(\mathbb{T})$ – the space of the continuous functions on \mathbb{T} , in the place of $L_\infty(\mathbb{T})$. Best trigonometric approximation of a function $f \in L_p(\mathbb{T})$ is given by

$$E_n^T(f)_p = \inf_{\tau \in T_n} \|f - \tau\|_p,$$

where T_n denotes the set of the trigonometric polynomials of degree at most n and $\|\circ\|_p$ denotes the usual L_p -norm on \mathbb{T} .

The error $E_n^T(f)_p$ is estimated by the so-called classical moduli of smoothness. To recall, the modulus of smoothness of order $r \in \mathbb{N}$ of $f \in L_p(\mathbb{T})$ is defined by

$$\omega_r(f, t)_p = \sup_{0 < h \leq t} \|\Delta_h^r f\|_p,$$

where the centred finite difference of order $r \in \mathbb{N}$ of f is given by

$$\Delta_h^r f(x) = \sum_{k=0}^r (-1)^k \binom{r}{k} f(x + (r/2 - k)h).$$

*Supported by grant DDVU 02/30 of the Fund for Scientific Research of the Bulgarian Ministry of Education and Science.

The following relation between $E_n^T(f)_p$ and $\omega_r(f, t)_p$ is a classical result in approximation theory (see for example [4, Ch. 7] or [15, 5.1.32, 6.1.1])

$$\begin{aligned} E_n^T(f)_p &\leq c \omega_r(f, n^{-1})_p, \\ \omega_r(f, t)_p &\leq c t^r \sum_{0 \leq k \leq 1/t} (k+1)^{r-1} E_k^T(f)_p. \end{aligned} \quad (1.1)$$

Above and in what follows we denote by c positive constants, which do not depend on the functions in the relations, nor on $n \in \mathbb{N}$ or $0 < t \leq t_0$. For $1 < p < \infty$ these estimates were improved by Timan [16, 17] (or see [4, Ch. 7] and [15, 6.1.5]) in the stronger forms

$$n^{-r} \left\{ \sum_{k=0}^n (k+1)^{sr-1} E_k^T(f)_p^s \right\}^{1/s} \leq c \omega_r(f, n^{-1})_p, \quad (1.2)$$

and

$$\omega_r(f, t)_p \leq c t^r \left\{ \sum_{0 \leq k \leq 1/t} (k+1)^{\sigma r-1} E_k^T(f)_p^\sigma \right\}^{1/\sigma}. \quad (1.3)$$

where $s = \max\{p, 2\}$ and $\sigma = \min\{p, 2\}$. The improved Jackson estimate (1.2) is also called a sharp Jackson inequality (see [3]).

Our main goal is to establish analogues of Timan's inequalities with moduli of smoothness, which are equivalent to zero if the function f is a trigonometric polynomial of a certain degree, that is, moduli which are invariant on such trigonometric polynomials. We call them *trigonometric moduli*. Such estimates look natural especially with regard to the Jackson-type inequalities (1.1) and (1.2) and the invariance of best trigonometric approximation. The method we shall use reduces the new inequalities to the classical. On the one hand, this gives a simpler proof of the Jackson inequality given in [9, Theorem 1.1], but on the other hand, we believe, this approach can lead to estimates of best approximation by other systems as well as of the rate of approximation of linear processes by moduli possessing a corresponding natural invariance.

A sharp Jackson inequality in a very general setting for multivariate functions was established by Dai, Ditzian and Tikhonov [3] in terms of K -functionals. Many concrete sharp Jackson estimates are derived, among which about the univariate best algebraic approximation by the Ditzian-Totik modulus, about the multivariate best trigonometric approximation by the classical modulus, and about best approximation by spherical harmonic polynomials by the Ditzian modulus on the sphere [6].

The following two inequalities between the classical moduli of different order are closely related to (1.2) and (1.3):

$$t^r \left\{ \int_t^{t_0} \frac{\omega_{r+1}(f, u)_p^s}{u^{sr+1}} du \right\}^{1/s} \leq c \omega_r(f, t)_p \quad (1.4)$$

and

$$\omega_r(f, t)_p \leq c t^r \left\{ \int_t^{t_0} \frac{\omega_{r+1}(f, u)_p^\sigma}{u^{\sigma r+1}} du \right\}^{1/\sigma}, \quad (1.5)$$

where $0 < t \leq t_0$. The former was established in [3, (1.6)]; the latter is due to Timan [16] (see also Zygmund [18]) and is referred to as an improved or sharp Marchaud inequality (see [4, p. 49] and the references cited below). These inequalities were extended to quite general function spaces in [5, 7, 8]. Also, Dai, Ditzian and Tikhonov [3, Theorems 5.3 and 5.5] (see also [2]) established more general forms in terms of K -functionals. There the multivariate form of (1.4) was proved [3, (2.17)]. In the present note we verify the analogues of (1.4) and (1.5) for the trigonometric moduli. Let us mention that their basic properties were established in [9, 10, 11] — they are just similar to those of the classical moduli.

The contents of the paper are organised as follows. In the next section we define the above-mentioned trigonometric moduli. Then in Section 3 and 4 we establish the analogues of (1.2)-(1.3) and (1.4)-(1.5), respectively, in their terms.

2 Trigonometric moduli of smoothness

We shall consider two different types of moduli that are identically zero on the trigonometric polynomials up to a certain degree. The first one is based on a modification of the finite differences, whereas the second on a modification of the approximated function.

It was Babenko, Chernykh and Shevaldin [1] who first defined a modulus, which is zero on the trigonometric polynomials of degree $r - 1$. It is given by

$$\tilde{\omega}_r^T(f, t)_p = \sup_{0 < h \leq t} \|\tilde{\Delta}_{r,h} f\|_p,$$

as the modified finite differences $\tilde{\Delta}_{r,h}$ were introduced by Shevaldin [13] (see also [12]) and are defined by

$$\tilde{\Delta}_{r,h} f(x) = \Delta_{r-1,h} \cdots \Delta_{1,h} \Delta_h f(x), \quad (2.1)$$

where

$$\Delta_{j,h} f(x) = f(x+h) - 2 \cos jh \cdot f(x) + f(x-h), \quad j = 1, 2, \dots$$

We have $\tilde{\omega}_r^T(f, t)_p \equiv 0$ iff $f \in T_{r-1}$.

To define the other modulus, let the 2π -periodic function \mathbf{a} be given on $[-\pi, \pi]$ by

$$\mathbf{a}(x) = \frac{1}{2} |x| (2\pi - |x|) \quad (2.2)$$

and let for $j \in \mathbb{N}_0$ the bounded linear operator $\mathfrak{A}_j : L_p(\mathbb{T}) \rightarrow L_p(\mathbb{T})$, $1 \leq p \leq \infty$, be defined by

$$\mathfrak{A}_j f = f + j^2 \mathbf{a} * f.$$

Above, $*$ denotes the convolution on $L_1(\mathbb{T})$

$$f * g(x) = \frac{1}{2\pi} \int_{\mathbb{T}} f(x-y)g(y) dy.$$

Further, for $r \in \mathbb{N}_0$ we set

$$\mathfrak{F}_r = \mathfrak{A}_r \cdots \mathfrak{A}_0.$$

Now, we define the trigonometric modulus of smoothness

$$\omega_r^T(f, t)_p = \sup_{0 < h \leq t} \|\Delta_h^{2r-1} \mathfrak{F}_{r-1} f\|_p.$$

It has the property that $\omega_r^T(f, t)_p \equiv 0$ iff $f \in T_{r-1}$ as it was established in [9, 11].

In [9, 10, 11] it was shown that the moduli $\omega_r^T(f, t)_p$ and $\tilde{\omega}_r^T(f, t)_p$ characterize the rate of best trigonometric approximation just similarly as the classical modulus in any homogeneous Banach space of periodic functions and, in particular, in L_p . Babenko, Chernykh and Shevaldin [1] proved the Jackson estimate (1.1) in the case $p = 2$ for the modulus $\tilde{\omega}_r^T(f, t)_2$. Shevaldin [14] verified it for $p = \infty$ and $r = 2$.

Let us explicitly point out that except for the case $r = 1$ when both $\tilde{\Delta}_{r,h} f$ and $\Delta_h^{2r-1} \mathfrak{F}_{r-1} f$ coincide with the symmetric finite difference of first order, there does not exist a simple algebraic connection between them. For $r \geq 2$ the finite difference $\tilde{\Delta}_{r,h} f$ is a linear combination of the values of the function f at the nodes $x + (r - k - 1/2)h$, $k = 0, \dots, 2r - 1$, whose coefficients depend on h . Basic properties of these coefficients including their explicit form are given e.g. in [10, Section 2] (see also [13]). On the other hand, the finite difference $\Delta_h^{2r-1} \mathfrak{F}_{r-1} f$ is the classical symmetric finite difference on the same nodes but of the function $\mathfrak{F}_{r-1} f$.

We can draw another line of comparison between the two finite differences. The relation (2.1) implies by induction the following representation

$$\tilde{\Delta}_{r,h} f = \Delta_h^{2r-1} f + \sum_{k=1}^{r-1} \lambda_k(h) \Delta_h^{2(r-k)-1} f,$$

where $\lambda_k(h)$ are even trigonometric polynomials. Whereas in the other case, since Δ_h and the convolution commute, we have

$$\Delta_h^{2r-1} \mathfrak{F}_{r-1} f = \Delta_h^{2r-1} f + \mathfrak{c} * (\Delta_h^{2r-1} f)$$

with an appropriate even 2π -periodic continuous function \mathfrak{c} (see [11, (1.10)]).

However, the moduli $\tilde{\omega}_r^T(f, t)_p$ and $\omega_r^T(f, t)_p$ are equivalent in the sense that there exists a positive constant c , whose value is independent of f and $t \leq t_0$, such that

$$c^{-1} \omega_r^T(f, t)_p \leq \tilde{\omega}_r^T(f, t)_p \leq c \omega_r^T(f, t)_p,$$

as it follows from results in [9, 10, 11] (see the end of the proof of Theorem 3.1 below).

3 An improved Jackson inequality

We shall establish the analogues of (1.2) and (1.3) for the trigonometric moduli $\omega_r^T(f, t)_p$ and $\tilde{\omega}_r^T(f, t)_p$.

Theorem 3.1. *Let $f \in L_p(\mathbb{T})$, $1 < p < \infty$, $s = \max\{p, 2\}$, $\sigma = \min\{p, 2\}$ and $r \in \mathbb{N}$. Then*

$$n^{1-2r} \left\{ \sum_{k=r-1}^n (k+1)^{s(2r-1)-1} E_k^T(f)_p^s \right\}^{1/s} \leq c \omega_r^T(f, n^{-1})_p \quad (3.1)$$

and

$$\omega_r^T(f, t)_p \leq c t^{2r-1} \left\{ \sum_{r-1 \leq k \leq 1/t} (k+1)^{\sigma(2r-1)-1} E_k^T(f)_p^\sigma \right\}^{1/\sigma}, \quad 0 < t \leq 1/r. \quad (3.2)$$

The inequalities remain valid with $\tilde{\omega}_r^T(f, t)_p$ in the place of $\omega_r^T(f, t)_p$.

Proof. The moduli $\omega_1^T(f, t)_p$ and $\tilde{\omega}_1^T(f, t)_p$ coincide with $\omega_1(f, t)_p$. So it remains to prove the theorem for $r \geq 2$.

Let \mathfrak{b}_j , $j \in \mathbb{N}_0$, be 2π -periodic and defined on $[-\pi, \pi]$ by

$$\mathfrak{b}_j(x) = j(|x| - \pi) \sin |jx|$$

and let

$$\mathfrak{B}_j F = F + \mathfrak{b}_j * F.$$

In [11, Proposition 2.4] we showed that for the bounded operator $\mathfrak{E}_r = \mathfrak{B}_r \cdots \mathfrak{B}_0 : L_p(\mathbb{T}) \rightarrow L_p(\mathbb{T})$ we have

$$\mathfrak{E}_r \mathfrak{F}_r f = f - S_r f, \quad f \in L_p(\mathbb{T}),$$

where $S_r f$ is the r th partial sum of the Fourier series of f . Also, as it follows directly from their definition, both operators \mathfrak{F}_r and \mathfrak{E}_r map the set of trigonometric polynomials T_k into itself for any k . Consequently, we get for $k \geq r-1$ and τ_k the trigonometric polynomial of degree k of best L_p -approximation of $\mathfrak{F}_{r-1} f$ the relation

$$\begin{aligned} E_k^T(f)_p &= E_k^T(\mathfrak{E}_{r-1} \mathfrak{F}_{r-1} f)_p \leq \|\mathfrak{E}_{r-1} \mathfrak{F}_{r-1} f - \mathfrak{E}_{r-1} \tau_k\|_p \\ &\leq c \|\mathfrak{F}_{r-1} f - \tau_k\|_p = c E_k^T(\mathfrak{F}_{r-1} f)_p. \end{aligned}$$

Now, (1.2) with $\mathfrak{F}_{r-1} f$ in the place of f and $2r-1$ in the place of r directly implies (3.1).

Similarly, (1.3) and the estimate $E_k^T(\mathfrak{F}_{r-1} f)_p \leq c E_k^T(f)_p$, $k \in \mathbb{N}_0$, imply

$$\omega_r^T(f, t)_p \leq c t^{2r-1} \left\{ \sum_{0 \leq k \leq 1/t} (k+1)^{\sigma(2r-1)-1} E_k^T(f)_p^\sigma \right\}^{1/\sigma}.$$

Next, we split the sum on the right hand-side into two parts for $0 \leq k \leq r-2$ and $r-1 \leq k \leq 1/t$. We estimate above the summands of the first sum using that $k \leq r-1$ and $E_k^T(f)_p \leq \|f\|_p$ to get

$$\omega_r^T(f, t)_p \leq c t^{2r-1} \left\{ \sum_{r-1 \leq k \leq 1/t} (k+1)^{\sigma(2r-1)-1} E_k^T(f)_p^\sigma + r^{\sigma(2r-1)-1} \|f\|_p^\sigma \right\}^{1/\sigma}.$$

Now, we replace above f with $f - \tau_{r-1}$, where τ_{r-1} is the trigonometric polynomial of best L_p -approximation of f of degree $r-1$, and use the invariance of $\omega_r^T(f, t)_p$ and $E_k^T(f)_p$, $k \geq r-1$, under addition of trigonometric polynomials of that degree to arrive at (3.2).

The inequalities for $\tilde{\omega}_r^T(f, t)_p$ are derived immediately from those for $\omega_r^T(f, t)_p$ because both moduli are equivalent to the same K -functional, namely,

$$K_r^T(f, t)_p = \inf_{g \in W_p^{2r-1}(\mathbb{T})} \{ \|f - g\|_p + t^{2r-1} \|\tilde{D}_r g\|_p \},$$

where \tilde{D}_r is the differential operator whose kernel is T_{r-1} (see [10, Theorem 4.2] and [11, Theorem 2.1]). \square

4 Improved relations between trigonometric moduli of different order

As we mentioned in the Introduction, the trigonometric moduli possess properties just similar to those of the classical one. In addition, they satisfy the following sharpened forms of the inequality $\omega_{r+1}^T(f, t)_p \leq c \omega_r^T(f, t)_p$ and of the Marchaud inequality.

Theorem 4.1. *Let $f \in L_p(\mathbb{T})$, $1 < p < \infty$, $s = \max\{p, 2\}$, $\sigma = \min\{p, 2\}$ and $r \in \mathbb{N}$. Then for $0 < t \leq t_0$ there hold*

$$t^{2r-1} \left\{ \int_t^{t_0} \frac{\omega_{r+1}^T(f, u)_p^s}{u^{s(2r-1)+1}} du \right\}^{1/s} \leq c \omega_r^T(f, t)_p \quad (4.1)$$

and

$$\omega_r^T(f, t)_p \leq c t^{2r-1} \left[\left\{ \int_t^{t_0} \frac{\omega_{r+1}^T(f, u)_p^\sigma}{u^{\sigma(2r-1)+1}} du \right\}^{1/\sigma} + \|f\|_p \right]. \quad (4.2)$$

The inequalities remain valid with $\tilde{\omega}_r^T(f, t)_p$ in the place of $\omega_r^T(f, t)_p$.

Proof. Iterating [3, (1.6)] (or see (1.4)), we get the inequality

$$t^{2r-1} \left\{ \int_t^{t_0} \frac{\omega_{2r+1}(f, u)_p^s}{u^{s(2r-1)+1}} du \right\}^{1/s} \leq c \omega_{2r-1}(f, t)_p. \quad (4.3)$$

Set $F = \mathfrak{F}_{r-1}f$. Then $\mathfrak{F}_r f = F + r^2 \mathbf{a} * F$. In [11, (3.2)] it was proved that $(\mathbf{a} * g)'' = g + \text{const}$ for any $g \in L_p(\mathbb{T})$. Then, using basic properties of the classical modulus, we get

$$\omega_{r+1}^T(f, u)_p^s = \omega_{2r+1}(\mathfrak{F}_r f, u)_p^s \leq c [\omega_{2r+1}(F, u)_p^s + u^{2s} \omega_{2r-1}(F, u)_p^s]. \quad (4.4)$$

For the first term on the right above we get by (4.3) with f replaced by F

$$t^{2r-1} \left\{ \int_t^{t_0} \frac{\omega_{2r+1}(F, u)_p^s}{u^{s(2r-1)+1}} du \right\}^{1/s} \leq c \omega_r^T(f, t)_p. \quad (4.5)$$

To estimate the second term on the right of (4.4), we proceed as follows. Let $F_t \in W_p^{2r-1}(\mathbb{T})$ be such that

$$\|F - F_t\|_p \leq c \omega_{2r-1}(F, t)_p \quad (4.6)$$

and

$$t^{2r-1} \|F_t^{(2r-1)}\|_p \leq c \omega_{2r-1}(F, t)_p. \quad (4.7)$$

For F_t one can take the Steklov mean of F (see e.g. [4, p. 177]). Then we have by basic properties of the classical modulus

$$u^{2s} \omega_{2r-1}(F, u)_p^s \leq c \|F - F_t\|_p^s + u^{s(2r+1)} \|F_t^{(2r-1)}\|_p^s$$

where $0 < u \leq t_0$, and, consequently,

$$\begin{aligned} t^{2r-1} \left\{ \int_t^{t_0} \frac{u^{2s} \omega_{2r-1}(F, u)_p^s}{u^{s(2r-1)+1}} du \right\}^{1/s} \\ \leq c \|F - F_t\|_p + c t^{2r-1} \|F_t^{(2r-1)}\|_p \leq c \omega_r^T(f, t)_p, \end{aligned} \quad (4.8)$$

as at the last step we have applied (4.6)-(4.7) and $\omega_r^T(f, t)_p = \omega_{2r-1}(F, t)_p$.

Now, (4.4), (4.5) and (4.8) imply (4.1).

We proceed to the proof of (4.2). Iterating the sharp Marchaud inequality (1.5), we arrive at

$$\omega_{2r-1}(f, t)_p \leq c t^{2r-1} \left\{ \int_t^{t_0} \frac{\omega_{2r+1}(f, u)_p^\sigma}{u^{\sigma(2r-1)+1}} du \right\}^{1/\sigma}.$$

With $\mathfrak{F}_r f$ in the place of f it yields

$$\omega_{2r-1}(\mathfrak{F}_r f, t)_p \leq c t^{2r-1} \left\{ \int_t^{t_0} \frac{\omega_{r+1}^T(f, u)_p^\sigma}{u^{\sigma(2r-1)+1}} du \right\}^{1/\sigma}.$$

Thus it remains to show that

$$\omega_r^T(f, t)_p = \omega_{2r-1}(\mathfrak{F}_{r-1}f, t)_p \leq c (\omega_{2r-1}(\mathfrak{F}_r f, t)_p + t^{2r-1} \|f\|_p). \quad (4.9)$$

To verify the latter, we take into account that $\mathfrak{F}_r f = \mathfrak{A}_r F$ with $F = \mathfrak{F}_{r-1} f$; hence $\mathfrak{B}_r \mathfrak{F}_r f = \mathfrak{B}_r \mathfrak{A}_r F = F + \eta_r * F$ with $\eta_r(x) = -1 - 2 \cos rx$ as was established in [11, (2.9)]. Set $G = \mathfrak{F}_r f$ and let $G_t \in W_p^{2r-1}(\mathbb{T})$ satisfy (4.6)-(4.7) for G in the place of F . Then

$$\begin{aligned} \omega_{2r-1}(F, t)_p &\leq \omega_{2r-1}(\mathfrak{B}_r G - \mathfrak{B}_r G_t, t)_p + \omega_{2r-1}(G_t, t)_p \\ &\quad + \omega_{2r-1}(\mathfrak{b}_r * G_t, t)_p + \omega_{2r-1}(\eta_r * F, t)_p \\ &\leq c \|\mathfrak{B}_r G - \mathfrak{B}_r G_t\|_p + t^{2r-1} \|G_t^{(2r-1)}\|_p \\ &\quad + t^{2r-1} \|\mathfrak{b}_r * G_t^{(2r-1)}\|_p + t^{2r-1} \|\eta^{(2r-1)} * F\|_p \\ &\leq c(\|G - G_t\|_p + t^{2r-1} \|G_t^{(2r-1)}\|_p + t^{2r-1} \|F\|_p) \\ &\leq c(\omega_{2r-1}(G, t)_p + t^{2r-1} \|f\|_p). \end{aligned}$$

Thus (4.9) is proved. \square

Remark 4.2. The inequalities (4.1) and (4.2) can be verified by means of (3.1) and (3.2) (see the proof of [3, Theorem 5.3]) and vice versa (the proof of [4, Theorem 3.4, Ch. 7]). Moreover, such a proof maybe considered even simpler and shorter than the one we used here. However, we preferred to use an approach which is based on the properties of the classical moduli and is independent of the relation of the new moduli to an approximation process. It demonstrates the advantages of the connection between $\omega_r^T(f, t)_p$ and the classical moduli in transferring properties between them and can be applied to define moduli appropriate for other approximation operators and establish their properties.

Remark 4.3. Replacing in (4.2) f with $f - \tau_{r-1}$, where τ_{r-1} is the trigonometric polynomial of best L_p -approximation of f of degree $r-1$, we immediately arrive at its slightly stronger form

$$\omega_r^T(f, t)_p \leq c t^{2r-1} \left[\left\{ \int_t^{t_0} \frac{\omega_{r+1}^T(f, u)_p^\sigma}{u^{\sigma(2r-1)+1}} du \right\}^{1/\sigma} + E_{r-1}^T(f)_p \right].$$

Acknowledgments. The research was conducted during my stay in Centre de Recerca Matemàtica, Bellaterra, Barcelona in February 2012 on the programme *Approximation Theory and Fourier Analysis*. I am especially thankful to Prof. Sergey Tikhonov for posing the problem and discussions on the presented results. I am also thankful to the referee whose remarks improved the exposition.

References

- [1] A.G. Babenko, N.I. Chernykh, V.T. Shevaldin, “The Jackson-Stechkin inequality in L_2 with a trigonometric modulus of continuity,” *Mat. Zametki*, **65**(6), 928–932 (1999) [in Russian]; English translation: *Math. Notes*, **65**(5-6), 777–781 (1999).

- [2] F. Dai, Z. Ditzian, “Littlewood-Paley theory and sharp Marchaud inequality,” *Acta Sci. Math. (Széged)*, **71**, 65–90 (2005).
- [3] F. Dai, Z. Ditzian, S. Tikhonov, “Sharp Jackson inequalities,” *J. Approx. Theory*, **151**, 86–112 (2008).
- [4] R.A. DeVore, G.G. Lorentz, *Constructive Approximation*, Springer Verlag, Berlin (1993).
- [5] Z. Ditzian, “On the Marchaud-type inequality,” *Proc. Amer. Math. Soc.*, **103**, 198–202 (1988).
- [6] Z. Ditzian, “A modulus of smoothness on the sphere,” *J. D’Analyse Math.*, **79**, 189–200 (1999).
- [7] Z. Ditzian, A. Prymak, “Sharp Marchaud and converse inequalities in Orlicz spaces,” *Proc. Amer. Math. Soc.*, **135**, 1115–1121 (2007).
- [8] Z. Ditzian, A. Prymak, “Extension technique and estimates for moduli of smoothness on domains in \mathbb{R}^d ,” *East J. Approx.*, **17**(2), 171–179 (2011).
- [9] B.R. Draganov, “A new modulus of smoothness for trigonometric polynomial approximation,” *East J. Approx.*, **8**(4), 465–499 (2002).
- [10] B.R. Draganov, P.E. Parvanov, “On estimating the rate of best trigonometric approximation by a modulus of smoothness,” *Acta Math. Hungar.*, **131**, 360–379 (2011).
- [11] B.R. Draganov, “Estimating the rate of best trigonometric approximation in homogeneous Banach spaces by moduli of smoothness,” Proc. intern. conf. “Constructive Theory of Functions”, Sozopol 2010: In Memory of Borislav Bojanov, Eds. G. Nikolov and R. Uluchev, Marin Drinov Academic Publishing House, Sofia, 80–92 (2012).
- [12] A. Sharma, I. Tzimbalario, “Some linear differential operators and generalized finite differences,” *Mat. Zametki* **21**(2), 161–172 (1977) [in Russian]; English translation: *Math. Notes* **21**(1-2), 91–97 (1977).
- [13] V.T. Shevaldin, “Extremal interpolation with smallest value of the norm of a linear differential operator,” *Mat. Zametki* **27**(5), 721–740 (1980) [in Russian]; English translation: *Math. Notes* **27**(5-6), 344–354 (1980).
- [14] V.T. Shevaldin, “The Jackson-Stechkin inequality in the space $C(\mathbb{T})$ with trigonometric continuity modulus annihilating the first harmonics,” *Proc. Steklov Inst. Math., Approximation Theory, Asymptotic expansions, suppl. 1*, 206–213 (2001).
- [15] A.F. Timan, *Theory of Approximation of Functions of a Real Variable*, Fizmatgiz, Moscow (1960); English translation: Pergamon Press, Macmillan, New York (1963).

- [16] M.F. Timan, “Converse theorems of the constructive theory of functions in spaces L_p ,” *Mat. Sb.*, **46**(88), 125–132 (1958) [in Russian].
- [17] M.F. Timan, “On Jackson’s theorem in L_p spaces,” *Ukrain. Math. Zh.*, **18**, 134–137 (1966) [in Russian].
- [18] A. Zygmund, “A remark on the integral modulus of continuity,” *Univ. Nac. Tucuman Rev. Ser. A*, **7**, 259–269 (1950).

Faculty of Mathematics and Informatics
University of Sofia
5 James Bourchier Blvd.
1164 Sofia, Bulgaria

Institute of Mathematics and Informatics
Bulgarian Academy of Sciences
bl. 8 Acad. G. Bonchev Str.
1113 Sofia, Bulgaria

`bdraganov@fmi.uni-sofia.bg`